

**AD-755 175**

# **Biological Sonar Systems**

**a bionics survey**

**Texas University**

**prepared for**

**Naval Ship Systems Command**

**SEPTEMBER 1972**

Distributed By:

**NTIS**

**National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE**

AD755175

# APPLIED RESEARCH LABORATORIES

ARL-TR-72-34  
5 September 1972

Copy No. 24

BIOLOGICAL SONAR SYSTEMS: A BIONICS SURVEY  
FINAL REPORT UNDER CONTRACT N00024-68-C-1339, ITEM 25  
1 November 1971 - 31 August 1972

K. Jerome Diercks

NAVAL SHIP SYSTEMS COMMAND  
Contract N00024-68-C-1339,  
Proj. Ser. No. S2616, Task 12867



Reproduced by  
NATIONAL TECHNICAL  
INFORMATION SERVICE  
U.S. Department of Commerce  
Springfield VA 22151

Approved for public release,  
distribution unlimited

172

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Applied Research Laboratories The University of Texas at Austin Austin, Texas 78712		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED ---	
3. REPORT TITLE BIOLOGICAL SONAR SYSTEMS: A BIONICS SURVEY FINAL REPORT UNDER CONTRACT NOOC24-68-C-1339, LINE ITEM 25			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) final report (1 November 1971 - 31 August 1972)			
5. AUTHOR(S) (First name, middle initial, last name) K. Jerome Diercks			
6. REPORT DATE 5 September 1972		7a. TOTAL NO. OF PAGES 172	7b. NO. OF REFS 175
8a. CONTRACT OR GRANT NO N00024-68-C-1339		9a. ORIGINATOR'S REPORT NUMBER(S) ARL-TR-72-34	
b. PROJECT NO S2616, Task 12867		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) ---	
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES ---		12. SPONSORING MILITARY ACTIVITY Naval Ship Systems Command Department of the Navy Washington, D. C. 20360	
13. ABSTRACT  The literature on biological sonar systems (bats, birds, marine mammals) was reviewed and past and present investigators were interviewed to ascertain the contributions, present and potential, of work in this field to high resolution sonar technology. It was concluded that there have been no contributions to date. Five areas of research were identified as potentially contributory: passive target ranging by the owl, signal design for target recognition, neural processing for target detection and recognition, the psychophysiology of sound localization, and the mechanics of signal generation by the small whales. The available system(s) and performance data on the bats and small whales are tabulated, and up-to-date bibliographies on each biological order are included.(U)			

UNCLASSIFIED

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Biological sonar systems						
Echolocation						
Bionics						
Bats						
Birds						
Whales						
Dolphins						
Porpoises						

ARL-TR-72-34  
5 September 1972

**BIOLOGICAL SONAR SYSTEMS: A BIONICS SURVEY**  
**FINAL REPORT UNDER CONTRACT N00024-68-C-1339, ITEM 25**  
1 November 1971 - 31 August 1972

K. Jerome Diercks

NAVAL SHIP SYSTEMS COMMAND  
Contract N00024-68-C-1339,  
Proj. Ser. No. S2616, Task 12867

Approved for public release;  
distribution unlimited.

**APPLIED RESEARCH LABORATORIES**  
**THE UNIVERSITY OF TEXAS AT AUSTIN**  
AUSTIN, TEXAS 78712

## TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	iii
I. INTRODUCTION	1
II. TARGET DETECTION AND CLASSIFICATION PHENOMENA	3
III. SIGNAL PROCESSING FOR DETECTION AND CLASSIFICATION	15
IV. MECHANISMS FOR SOUND GENERATION AND RECEPTION	21
V. PASSIVE/ACTIVE SONAR UTILIZATION	27
VI. CONCLUSIONS	29
REFERENCES	33
APPENDIX I	47
APPENDIX II	55
APPENDIX III	107
APPENDIX IV	123
APPENDIX V	151
APPENDIX VI	175

## I. INTRODUCTION

Applied Research Laboratories (ARL), The University of Texas at Austin, under Contract N00024-68-C-1339 with Naval Ship Systems Command, has performed a survey of bionics research on animal sonar systems. Funding for this work began 1 November 1971 and terminated 31 August 1972. This report describes the nature and scope of this effort and summarizes the conclusions reached.

The objective of this effort was to identify and specify the features of biological sonar systems which can be exploited by sonar technology or the knowledge of which might influence or alter sonar design philosophy. The biological forms considered in this survey are the insect-eating bats (Order Chiroptera; Suborder Microchiroptera), whales, dolphins, and porpoises (Order Cetacea, Suborder Odontoceti), and a few species of birds.

The literature relating to biosonar systems was reviewed. Nearly 600 documents relating to echoranging, physiological mechanisms of sound generation and reception, morphology, and ecology (diet, habitat, etc.) were collected. Included are approximately 85 foreign language reports (44 in Russian, 24 in French, and 17 in German) which are being translated (45 completed to date). A tabulation of data for each bioform is included in the appendices. Bibliographies of the documentation on biosonar systems are also included.

Most active members of the biosonar community and many members of the sonar engineering community at ARL were interviewed to achieve a critique of bionics research. A listing of the biosonar membership

contacted is included as Appendix I. A standardized questionnaire was employed to guide the discussions, namely:

- (1) Has bionics research on animal sonar systems contributed to synthetic sonar design or design philosophy? In what ways?
- (2) What areas of animal sonar research appear most likely to provide guideline information for synthetic sonar design or design philosophy?
- (3) In what ways can sonar engineering aid bionics research on animal sonar systems to derive guideline information for sonar technology?

The answer to the first question was unanimously, no. The answer to the second question is the principal topic of this report. There were two answers to question 3. The first, explicitly stated, was to provide improved instrumentation (broader bandwidth, greater sensitivity) for detection, recording, and analysis of biosonar signals. The second, implied, was to communicate the interests and requirements of sonar technology to provide guidelines for productive biosonar research. Apparently, previous attempts to achieve the latter--whether initiated by the engineer or biologist--have been essentially unsuccessful.

There are several excellent, comprehensive reviews of biosonar phenomena (Refs. 4, 41, 44, 70, 91, 107, 120, 122, 131, 147, 164, 170, 175), and this report will not attempt to reiterate these. Five broad categories of bionics information are identified: target detection phenomena, target classification phenomena, signal processing for detection and classification, passive/active sonar utilization, and mechanisms for sound generation and reception. The first two will be considered together.



## II. TARGET DETECTION AND CLASSIFICATION PHENOMENA

Sonar technology stands in awe of the apparent ease with which biosonar systems detect and classify targets. In many instances the operating sonar/target parameter relationships significantly exceed those achieved by synthetic means. For example, bats are able to detect and avoid wire obstacles in their flight path at wavelength-to-target diameter ( $\lambda/d$ ) ratio values of 15 to 25 (Refs. 48,49,53,63, 66,84,98,140,158). Food prey, like gnats and mosquitoes, yield ratio values of 1 to 10, depending upon the target dimension selected. The dolphin readily finds objects  $0.25$  to  $0.5\lambda$  in dimension (Refs. 74,78) and avoids obstacles  $0.1\lambda$  in dimension (Refs. 26,27,77,78,111). It is reported able to resolve\* targets separated 1 to 2 mm ( $0.05$  to  $0.1\lambda$ ) in range and  $0.25^\circ$  to  $1^\circ$  in azimuth (Ref. 5). The owl determines bearing and apparently range passively and suffers but slightly doing so with only one receiver (ear) (Ref. 79,125).

Members of both biological orders have been trained to discriminate between 2- and 3-dimensional targets of differing shapes, forms, and materials (Refs. 6,17,38,54,67,78,81,83,145,146,152,168). Once the animal comprehends the task requirement, identification of a positive (versus any other) target by echolocation (Refs. 81,83,128,135), discrimination performance improves relatively rapidly. That is, the animal readily learns (identifies) characteristics of the target echo that identify the positive target (and in absentia, also identify any negative target). It has been necessary to "teach" some dolphins to echolocate in a testing environment (Refs. 3,16,109), but once "taught" they perform (discriminate between targets) as well as any other so tested. The limits of ability to discriminate size (acoustic cross

---

\* It is probable that the animal is detecting qualitative differences between the echoes from the test targets and a single target presented simultaneously without actually measuring (resolving) the separation between the test targets.

section, or target strength) have been determined for several species (Refs. 3,17,41,128,157). A working value is 10% difference in size (Refs. 3,113,152) or 1 dB difference in target strength (Refs. 41,128).

The detection/avoidance phenomena imply apparent operation in a low noise background (at least within the requisite bandwidth, whatever that might be), or the ability to process the echo signals to achieve an effectively high signal level (or low noise level), or probably both. Attempts to mask echolocation by projecting high intensity, broadband noise into the flight path of bats have been only partially successful (Refs. 50,53,60). It appears there is a physiological signal-to-noise ratio greatly exceeding the physical value, and knowledge of how this is accomplished would assuredly benefit sonar technology. Current efforts in electrophysiological biosonar measurements are rapidly increasing our understanding of the mechanisms involved and should be supported (see next section).

A "figure-of-merit" (Ref. 44) calculated for a post World War II search radar, for two species of FM bats, and for a dolphin yields comparable performance values for the bats and the radar (Ref. 44), but indicates the dolphin is several orders of magnitude poorer (Ref. 137). However, the latter result is questionable due to poor validity of the measurements assumed in performing the calculation.

The owl's ability to localize and range passively, insofar as it is documented and understood, is of prime interest to the sonar community, and increased support of efforts aimed at understanding this ability is highly recommended.

In most parameters, synthetic sonars excel their biosonar counterparts. The exceptions are in bandwidth and signal-to-noise ratio. Recent developments in sonar technology indicate a capability of achieving comparable bandwidths (Refs. 14,34). Detection and

classification, then, seem not to be a question of capability, but of ability. A design philosophy which specifies mechanical processing of echoes precludes a legitimate comparison with biological systems. They are not comparable at this time. Sonar technology has achieved only marginal success with wholly mechanical systems, regardless of their capabilities. Knowledge of biosonar capabilities and abilities thus begs the recommendation that the ear-brain processor (the sonar operator) be reintegrated into sonar evaluation philosophy.

Little is known about echolocation by whales in the wild. They are intelligent and curious, and the observer's presence assuredly alters their behavior and stimulates acoustic activity probably more excitatory than echolocational in nature. Bats, however, seem little bothered by the presence of man in their environs and may be observed without disturbing their normal behavior. Some bats capture as many as 15 to 25 insects per minute under natural conditions (Refs. 51,118), but in the laboratory must be trained to discriminate (i.e., be provided experience with) non-food objects. They will attack any object from pebble size to basketball size the first few times it is thrown into their flight path (Ref. 62). (After training, especially when satiated, bats will often attack these non-food objects, apparently for diversion (Ref. 164).) A dolphin that has been trained to discriminate artificial (non-food) targets and has been fed whole dead fish, which it sometimes must retrieve acoustically, will ignore a similar specimen when presented as a target object in a testing environment (Ref. 10). Thus, it appears that routine classification is "monopolar" and intrinsic to the context in which it is performed. For instance, an echo indicates a likely food object, or positive target, and therefore initiates pursuit or approach, or it doesn't. In the wild echolocating animals, such as bats, may actively classify little more than surface, whether it is land or water (Refs. 46,121). The insectivorous bat in a hunting context pursues any object in translational motion. (It does not attack rustling leaves, and it

soon learns that not all "flying" objects are food.) The dolphin learns to recognize, and to respond to, an echo signature in context and to ignore it out of context.


This consideration suggests a fundamental guideline for bionics research, that is, to vary the animal's sonar problem and monitor the development of its solution rather than to evaluate the apparent solution after it has developed, which is the usual practice. This will allow description of the testing procedure (the signal design) employed by the animal to achieve a solution. Then, with knowledge of the target(s), it would be possible to provide a description of the problem and, thus, its solution. The merit of this procedure is universally recognized, but it has not been practiced for perhaps obvious reasons. An operational criterion for inferring the existence of solution is the stabilization of behavior. It is then a simple matter to record a short sequence of measurements after this criterion is reached. The alternative is to record many measurements over an indeterminate period during development of the solution, and the cost may be excessive. However, the former approach yields no way of knowing if the apparent solution is optimal, a consequence of adaptive testing within the constraints of the animal's capabilities, or is merely effective, a manifestation of the only overt solution the animal is capable of, regardless of the problem. From the viewpoint of sonar technology, the potential return from a "developmental" approach would seem to more than offset its probable cost.

A second alternative is to continually vary the sonar problem, by substitution or replacement of targets, once it has been established that the animal understands what it is required to do, and to monitor the changes in the animal's echolocation behavior, which are then correlated with the assumed changes in the problem (Ref. 146). The dangers of this procedure are that the animal may fail to understand the behavior required of it and cease to perform, or that it may


proceed with the overt behavior of the testing paradigm without performing any acoustic discriminations (Ref. 10). Any conclusions drawn from this latter event would be most misleading. However, use of this technique has illuminated apparent differences in the signals employed by one species of FM bat for "orientation," "target location," and "classification" (Refs. 146,152).

The small whales use a transient-like, usually ultrasonic click for echolocation (Refs. 12,13,41). This is a broad bandwidth signal: 3 to 5 octaves,  $Q \approx 1$ . The generic form is illustrated in Dwg. AS-72-923. The mechanism of generation is moot (see Section IV on mechanisms of sound generation). The signal is directed forward in a narrow, conical beam ( $20^\circ$  to  $90^\circ$  beamwidth, depending on frequency) (Refs. 39-41,112,138,139,170). Signal frequency decreases with angle from the acoustic axis (Refs. 34,112), a feature which may be exploited for target localization and/or classification. The dolphins scan the forward volume (direction of motion) by rotating the head through an elliptical path (up-and-down as well as side-to-side motion) (Refs. 41, 111,112,128,170). It is not known whether this is related to transmission, reception, or both.

Whales employ a somewhat stereotyped transmission format for echolocation of artificial targets in a tank (Refs. 13,38,41,74, 101,113,128). As the animal closes range on the target(s), it transmits usually at an increasing rate and reduces signal intensity. There is no ambiguity; the next transmission occurs only after receipt of the echo from the preceding transmission (Ref. 101). The generator may, in fact, be triggered by feedback from the receiver, although experienced test animals seem able to consciously control repetition rate, transmitting only often enough to maintain tracking and/or to achieve target identification ( $\sim 25$  to  $50$  per second) (Ref. 10). Signal intensity is reduced, apparently to minimize extraneous returns from other targets or surfaces. The dynamic range of intensities over



WAVEFORM



SPECTRUM

# GENERIC DOLPHIN ECHOLOCATION SIGNAL

a 3 to 5 m test path is greater than 35 dB (exceeds the range of available recording instruments). No reliable spectral changes have been observed over this distance (Refs. 12,13,135).

Bats use a diversity of signal forms for echolocation: short duration (1 to 5 msec) frequency modulated (FM) slides (Refs. 48,61,65,118,130,145,146,149,161,168); long duration (20 to 100 msec), constant frequency (CF) bursts, terminated with a short duration FM slide (Refs. 1,61,65,85,116,141,142,172); a mixed form consisting of a short duration CF burst terminated with a similar duration FM slide (Refs. 61,65,117-119,124,157); and short duration (~1 msec) CF pulses (Refs. 48,98). (One species of Megachiropteran bat, Rousettus aegypticus, generates an audible transient by clicking its tongue (Refs. 49,87).) Signals may be pure (Refs. 48,65,97,130,147,149) or contain one or more higher harmonics (Refs. 48,61,65,116,118,130,140,142), one of which may dominate (Refs. 61,65,118,130,140,142). FM bandwidths range from about 0.2 to 2 octave, depending upon species (Refs. 48,116,118,130,145,146,149,173), signal form (Refs. 61,65,85,130), and phase of pursuit/capture or obstacle avoidance maneuver (Refs. 145,146,157,158,168). Sounds are generated in the larynx (see Section IV on mechanisms of sound generation). The FM bats radiate through the open mouth, which is formed like a megaphone. The directivity pattern is broad (~45° to 90° at the half-power points)(Refs. 92,148,155), but the sound field is frequency structured and it is thought that bats achieve fine spatial resolution by exploiting this feature (Ref. 64). They do not appear to scan mechanically during pursuit/tracking (Ref. 167).

The CF bats in general radiate through the nostrils, which are separated by one-half wavelength at the dominant frequency and may be enclosed in a tissue appendage (nose leaf, horseshoe) which aids in directing the transmission (Refs. 44,92,95,97,98,100,131,154,172).

Directivity is greater than for FM bats (beamwidths about  $30^\circ$  to  $60^\circ$  at the half-power points)(Refs. 92,97,100,141,148,154,156), and there is noticeable mechanical scanning of the head during pursuit/tracking (Refs. 69,94,95,97,100,142).

Bats also employ a stereotyped transmission format. When searching they pulse at a low repetition rate (5 to 10 per second). Analogous to synthetic sonar usage, this may indicate an ability to detect at much longer ranges than ordinarily attributed to these animals (Ref. 46) or it may be merely a manifestation of energy conservation (Ref. 44). During pursuit/capture or obstacle avoidance, the repetition rate increases and signal intensity is reduced (Refs. 6,28,44,45,51,69,82,85,117,119,124,140,141,145,146,157,161). It is moot whether the onset of increased pulse repetition indicates the time of detection, classification, or merely initiation of pursuit. It is not unlikely that detection/classification occurs during search, perhaps aided by passive clues (Refs. 6,82,162) and possibly vision (Refs. 1,2,22,29,102,159). The FM bats initiate pursuit or avoidance at about 1 to 3 m distance from the target (Refs. 6,21,51,54,63,119,157,158), and the CF bats at about 3 to 10 m distance (Refs. 97,99,124).

The FM bats shorten the duration of their signals during pursuit/avoidance to prevent pulse-echo overlap, except in the final moments of capture, and they reduce the frequency range of their slides by successively omitting the higher frequency components (Refs. 28,45,51,145,157,168). Thus, resolution is progressively degraded (directivity and bandwidth are reduced) during pursuit/capture. Some species emit pulses in doublets during the later stages of pursuit/avoidance (Refs. 98,161). The intent of this change in transmission format is unknown.

Bats transmitting a mixed signal form follow the same protocol, except the signals are first shortened by eliminating the CF portion, then proceed as do the pure FM species (Ref. 157).



The CF bats permit pulse-echo overlap throughout the pursuit/avoidance maneuvers (Refs. 117,122,124,141). Some species shorten signal duration during target closure to maintain constant pulse-echo overlap (Refs. 117,119,122). The benefit derived from this is unknown.\* Other species initially increase the duration of their transmissions to increase pulse-echo overlap, then proceed by shortening the duration more or less linearly with range to keep the FM ending of the echo constant in time relative to the onset of the transmission (Refs. 117,122). The lengthy overlap probably permits accurate determination of the Doppler shift in echo frequency resulting from platform (bat) motion.\*\* The benefit derived from initially increasing overlap and then maintaining the relationship described is, again, unknown. Initial pulse-echo overlap may index detection. Increasing the overlap might then facilitate the initiation of pursuit at the earliest time (greatest target distance). The Rhinolophidae, the horseshoe bats, emit a long CF pulse during search, but change to a short pulse burst transmission mode when approaching an obstacle (Refs. 45,52,85,140-142). Both the search and burst pulses terminate with an FM sweep (Refs. 45,85,141,142). It is speculated the intent of this change in signal format is to increase the rate of information provided by the terminal FM portion of the echo. This interpretation seems overly simple. The total duration of a pulse burst is usually longer than a single long pulse (Refs. 85,140-142); pulse echo overlap is thus increased with no apparent synchrony between pulses and echoes. The bat may be able to attend to only the FM portions of the signals and thereby achieve some resolution. But then why does it emit the CF portion of the signal

---

It is possible the overlap generates a difference-frequency component which signals the generator to stop transmitting; i.e., the duration of overlap is a measure of the system's reaction time. The bat senses target closure by apperception of the shortening of signal duration.

\*\* Sensing of constancy of frequency shift may be apperceived as normal progression of target closure.

at all in this mode? These bats excel the FM species in detecting smaller obstacles at longer ranges (Refs. 84,85,124,140,141); accordingly, their sonar merits additional study.

There are no outstanding singular differences between the sonar problems confronting FM and CF bats, as manifested by their hunting behavior(s), diet(s), and ability to resolve artificial targets in the laboratory (Refs. 46,49,84,116,120,158,163). Both feed on flying insects; the CF bats, being somewhat larger in size than the FM bats, often attack larger species (Refs. 44,163,164). The CF bats are less agile flyers than some of the FM species and are able to hover in flight, which allows them to capture crawling or stationary prey (Refs. 98,120,162,163). The insectivorous bats, confronted with the problem of tracking small moving targets in an uncluttered environment, emit high intensity signals (SPL  $\sim 135$  dB re  $0.0002$  dyne/cm<sup>2</sup> at 10 cm), while the hovering bats, seeking large or motionless prey in oftentimes dense foliage, emit relatively low intensity signals (SPL  $\sim 75$  dB re  $0.0002$  dyne/cm<sup>2</sup> at 10 cm) (Ref. 43). The fishing bat, Noctilio leporinus, which is sometimes insectivorous, emits a relatively intense signal (Ref. 48). Many of the CF bats achieve a broad signal bandwidth by emitting many harmonics, but in an ill-defined format (Refs. 48,118).

Noctilio using a mixed signal form is able to detect small perturbations of the water surface to locate fish swimming just underneath, which it then gaffs with its feet (Refs. 20,120,157). It is able to detect small wires protruding above a smooth water surface (Ref. 157), as well as the "wake" of a fish traversing a wind rippled surface (Refs. 20,44,120). During pursuit/capture it shortens its signal by successively eliminating the CF portion to prevent pulse-echo overlap (Ref. 157). It employs a different transmission format to detect and avoid wire obstacles in its flight path

than it does to pursue fish. That is, it continues to transmit a mixed signal form throughout without shortening the pulse and uses pulse-echo overlap as it closes the target/obstacle (Ref. 158).

The megachiropteran bat, Rousettus sp., together with the South American Oilbird, Steatornis, and certain species of Asian Cave Swiftlets, Collocalia, emits click-like echolocation signals generated with the tongue and radiated from the corners of the mouth (Refs. 45, 44, 49, 55, 87, 115). All utilize the capability functionally, in lieu of vision, for navigation or orientation in the dark; none apparently uses it for detection and pursuit of prey (Refs. 43, 44, 49, 96, 102).

### III. SIGNAL PROCESSING FOR DETECTION AND CLASSIFICATION

Available evidence indicates that an animal's signal characteristics are matched to its receiver capabilities (hearing) (Refs. 5,30,41,57,61,65,68,70,75,86,103,162). Hearing sensitivity changes as the animal matures, and it apparently adapts its signal form to coincide with these changes (Ref. 65). After maturation there is evident adaptation to changes in sonar environments and/or tasks (Refs. 1,2,21,42,44,98,145,158,167).

There are few data relating directly to signal processing for target classification. The reason for this is principally procedural. Even when it is possible to monitor both the transmitted and target echo signals, as well as the animal's behavior, it is operationally difficult to ascertain the time of occurrence of, or to infer the analytic sequence leading to, a classification decision. There is sufficient signal-to-signal variation to generate quite misleading or erroneous conclusions should the improper moment or sequence be selected. Bat and dolphin signal forms have been mechanically simulated to examine echo characteristics of targets used in animal tests (Refs. 17,21,31,32,41,54,136,168,169). A study of target shape discrimination by the bat, Vampyrum spectrum, an FM bat, followed by analysis of target echoes generated by artificial signals, indicated learned discrimination based upon overall echo amplitude differences by one bat and upon frequency response differences by a second bat (Ref. 21). Other similar efforts have been noticeably less successful (Refs. 17,31,32,54,136,168,169). Correlations between identified echo signal characteristics and known target characteristics or between signal characteristics and the animal's discrimination performance are minimal and unreliable. Presentation of the data,

translated in frequency to human listeners, results in facile discrimination of perceived signal characteristics that identify a class of targets (e.g., "hard," "soft," spherical, cylindrical) (Refs. 32,168). Attempts to extract these clues by machine processing have been only marginally successful (Ref. 12,31,32).

Analyses of signals emitted by dolphins during target shape and/or material discriminations have been generally unrevealing; the signals (emitted and/or reflected) examined do not exhibit significant differences in form or spectrum (Refs. 12,13,135). There is a reported correlation between discrimination performance and echo structure (of elastic targets) (Ref. 35), but this is not unexpected. Finally, it has been speculated that bats and dolphins utilize learned target motion behavior (track, range rate) as a (perhaps the) classification clue (Ref. 56).

There are abundant data relating to signal processing for target/echo detection and localization. The results to date are both exciting and encouraging. Electrophysiological measurements of activity in peripheral and higher nervous centers in response to artificially and self-generated sounds have revealed unique capabilities related to the echolocational process(es) of both bats (Refs. 6,57,58, 59,61,64,65,69,70,103,106,127,162) and dolphins (Refs. 23,24,88). Work in this area is expanding. By 1973 the recording of radio-telemetered data from free-flying bats with microelectrodes chronically implanted in the auditory nervous system should be accomplished (Ref. 71). Progress on analogous measurements on dolphins is hindered principally by state-of-the-art knowledge of animal maintenance and care.

The auditory systems of echolocating bats are sensitive to ultrasonic frequencies, maximally so in the region(s) of their emitted sounds (Refs. 5,30,57,61,65,103,106,127,162). Also, they are capable of fast time resolution, and some species show an increased responsiveness to the second of a pair of sounds (an "echo" after

transmission) (Refs. 58,64,65). (Recent work on dolphins shows similar capabilities for those animals (Refs. 23,24).) Hearing sensitivity in those bats that transmit a relatively long duration, constant frequency (CF) signal is sharply tuned (effective roll-offs of several hundred decibels per octave change in frequency) to the dominant frequency emitted while the bat is at rest (Refs. 61,65,103,106,127,151). In flight these bats lower their transmitted signal frequency to compensate for Doppler shifts in the echo caused by source and/or target motion and thus maintain the echo at the frequency of maximum hearing sensitivity (Refs. 127,141-143,151). This has the effect of reducing apparent echo masking during pulse-echo overlap (Refs. 103,106,127,143) and of sharpening target localization by rejection of off-axis target echoes, which will be shifted less in frequency (proportional to the cosine of their angle with the acoustic axis) than those on-axis (Refs. 103,143). Apperception of the shift in signal frequency would yield a measure of range rate.

All species, CF and FM, reduce signal intensity during target closure, probably to minimize echo clutter (volume reverberation) from extraneous targets (Refs. 44,69,82,85,140,141,145,157).

The CF bats radiate sound from a dipole source\* (the nostrils, separated one-half wavelength at their most sensitive frequency) and scan the sound beam by moving the head during echo ranging (Refs. 97,98,142). They also move their pinnae (external ears) through a forward-to-side-looking arc while echolocating (Refs. 52,70,97,131-133,142). The pinnae alternate, one directed forward while the other is directed laterally. In the Greater Horseshoe bat, Rhinolophus ferrum equinum, the alternations occur in synchrony with the emitted pulses (Ref. 52). In other CF bats, the emission rate and ear movements are asynchronous (Refs. 69,70,133,142). The two scanning behaviors, projecting and receiving, together provide for crossrange (or azimuthal) tracking;

---

\* Except Chilonycteris p. parnellii which emits through the open mouth.

the latter alone may provide for crossrange rate determination. It has also been speculated that the ear movements, by providing a Doppler scan across frequency, may facilitate accurate frequency tracking (Ref. 151). Also, by orienting one receiver toward the target and the other orthogonally to this direction, the animal enhances its ability to detect the target echo in a noisy environment (Ref. 60).

The short FM slide (~10 kHz) terminating each CF transmission may function as an alerting signal during searching behavior when the animal is at rest. If, when searching, the bat transmits at or near its frequency of maximum sensitivity, the CF portion of an echo will be shifted upward to a less sensitive frequency region. However, the FM part of the echo will sweep down through the region of maximum sensitivity and thus alert the bat to the presence of a target. The utility of this function would be vitiated in flight, unless target motion shifted the echo frequency above the region of maximum sensitivity.

There has been much recent speculation about the intent and use of the CF and FM portions of the signals emitted by the "CF" bats, and, in comparison with the "pure FM" species, of the capabilities inherent in the two generic signal forms (Ref. 174). It is conceded the FM bats use an optimally Doppler tolerant waveform for determining target range (Refs. 7,174). The CF bats presumably minimize the effects of Doppler shifts in echo frequency upon target range determination by compensating for these shifts during transmission of the CF portion of the signal, i.e., by sliding the signal up or down in frequency to compensate for Doppler. The FM portion of the echo is maintained at the same "frequency" as that of the transmission to yield minimum ranging error (Ref. 174).

Hearing sensitivity in the FM bats is broadly tuned to cover the range of frequency changes and in some instances is bimodal, showing increased sensitivity over the ranges of the first and second harmonics of the transmission (Refs. 5,30,57,61,65,162). These bats continuously

reduce signal intensity during closure, presumably to minimize echo clutter from extraneous targets. They also shorten signal duration, to prevent pulse-echo overlap, and reduce FM bandwidth to about 15% of its initial (search) value during the pursuit/capture or obstacle avoidance maneuver (Refs. 45,51,54,145,146,168). Studies with two species of FM bats, the big-brown bat, Eptesicus fuscus, and the neotropical spear-nosed bat, Phyllostomus hastatus, have demonstrated the apparent use of correlational processing for range determination (Refs. 147,149). The test problem was seemingly unnatural for the bats; they were required to ascertain the range difference between fixed targets from a stationary perch. It is moot whether they rely on correlational processing during pursuit. The change in signal parameter values would continuously degrade the processing gain and resolution available, but this might be compensated for by the improvement in signal-to-noise ratio which accrues during closure.

Theoretical speculations on biological sonar phenomena are difficult to test; most are heuristic. The positive results obtained in the previously mentioned studies with Eptesicus and Phyllostomus are exceptional. More commonly, the applicability of a theory is called to question by demonstration of failure of occurrence of a requisite feature or event (Refs. 28,60,74,90,114,144,160) at the level of measurement. Claims of "solutions" to biological sonar problems (Refs. 7,8) are premature and pretentious; the problems have not yet been defined (Refs. 47,82,110,121,164-166), and the signal forms are not yet reliably known (Refs. 18,33,34). Such claims are merely statements of the capabilities and limitations of the signal form(s) examined, given the processing employed (Refs. 7,8,41); that performed by the animal is unknown but may be inferred from results like those obtained in the Eptesicus and Phyllostomus studies.



The projected sound field of an FM bat will be frequency structured, at least to the extent that frequency decreases with azimuthal angle. Receiving directivity is frequency dependent in a complex way (Refs. 6, 61, 64, 65), and it has been shown that, by sampling an echo at three (or more) frequencies within the swept band, bats could localize the source of an echo uniquely in the forward hemisphere (Ref. 64). It is possible the CF bats utilize the terminal FM portion of their signal similarly.

Attempts to derive possible target classification clues by machine processing of bat and dolphin signals, or bat-like and dolphin-like signal echoes from assorted test targets, have been disappointingly unrevealing (Refs. 12, 14, 31, 32). Presentation of the same signals, suitably translated in frequency to the audio range, to human listeners exposes obviously perceptible signal differences which are not revealed by the processing applied (Refs. 14, 31, 32). It is generally conceded that no extant machine or machine processing technique yields a meaningful analog of ear-brain processing (Refs. 25, 47, 121, 150).

#### IV. MECHANISMS FOR SOUND GENERATION AND RECEPTION

Bats generate sounds in their larynx,\* which is anatomically different from that of other mammals (Refs. 96,123), apparently evolved for production of the short duration, high intensity, ultrasonic signals used in echolocation. Ultrasonic capability is achieved by operating the vocal membranes at relatively high tension (Ref. 123). (See also, Ref. 173). Of significance in the present context is the ability to produce the high intensity levels recorded ( $\sim 110$  to  $130$  dB re  $0.0002 \text{ dyn/cm}^2$ , or  $35$  to  $45$  dB re  $1 \text{ } \mu\text{bar}$  at  $10 \text{ cm}^{**}$ ), given the minuscule size of the generating organ(s). This is apparently accomplished by causing a large overpressure at the distal port of the larynx; air is then metered through the laryngeal orifice for the duration and at a rate appropriate to the task (phase of the pursuit/avoidance maneuver) (Ref. 123). The transmission pattern is cyclic and correlates with respiration (Refs. 44,131,140,143,161).

The mechanism(s) of sound generation by whales is moot (Ref. 170). Three hypotheses prevail. First is that the sounds are generated in the larynx and transmitted via the musculature in the nasopharynx to the base of the skull, then along the maxillae (rostrum) into the water (Ref. 129). This hypothesis is not supported by anatomical or physiological evidence (Refs. 109,114). The other hypotheses are similar in the

---

\* The megachiropteran fruit bat, Rousettus, generates echolocation sounds by "clicking" its tongue.

\*\* Correcting for directivity yields source levels  $6$  to  $12$  dB lower than the values stated.

location of the generator, the nasal plugs in the bony nares,\* but differ in the mechanics of generation. In one theory sounds are generated by friction grating of the plugs against the opposing tissue; this is under muscular control (Refs. 37,39). The effect is either a staccato-like hammering of the skull, or cavitation as the plug moves, or both. The other hypothesis ascribes generation to flapping of the plug(s) against its opposing surface as air is metered past, much like a "bronx cheer" is generated with the lips (Refs. 108, 114). The effect here is, again, a hammering of the skull. Sounds are radiated via the fatty melon into the water, or along the maxillae into the water, or both.

The maxillae are apparently acoustically separated by cartilaginous tissue along most of their length (Ref. 37). Separation at the distal margin is nominally one-half wavelength at the peak signal frequency, implying a dipole source analogous to that described for the CF bats. Calculated directivity values for the bottlenose dolphin, Tursiops truncatus (Ref. 11), agree with measured values (Refs. 41,112), in support of the implied configuration of the radiator. Sound pressure levels ~110 to 130 dB re 1  $\mu$ bar at 1 m have been reported for several species (Ref. 41). (However, see Ref. 170).

The amount of available information in an echo is dependent upon the bandwidth of the signal interrogating the reflector. The bandwidth of a whale's signal (see Fig. 1) greatly exceeds that of synthetic sonars and, hence, yields potentially more information about the nature of a target. Recent efforts to exploit this knowledge by mechanically simulating a whale-like signal for high resolution target classification (Ref. 14), while successful, have also illuminated the gross inefficiencies of mechanical (electroacoustic) generation in comparison

---

\*This location has been confirmed by acoustical measurements performed on the animal (Ref. 33).

with its biological counterpart. It is difficult to achieve the desired bandwidth and intensity and directivity simultaneously.

Accordingly, the mechanism(s) of sound generation and radiation by the odontocete whales would seem to be a promising area for bionics research, providing inputs to the fields of acoustics and sonar engineering. Support in this area is recommended. The bats' phenomenal sound producing capabilities, while intriguing to both the biologist and acoustician, do not present any basis for exploitation by sonar technology at this time.

Behavioral and electrophysiological evidence reveal that the auditory sensitivities of the bats and whales are not exceptional (Refs. 9,30, 57,61,65,68,72,73,75,76,86,126,162,171). The external meatus (ear canal) of the whales appears to be vestigial and the tympanic membrane (ear drum) nonfunctional (Refs. 19,88). The path of sound reception in the dolphin, T. truncatus, is via the posterolateral portion of the mandible (lower jaw) to the auditory bulla (Refs. 23,41,88,108,109). Stimulation is effected by differential vibration of the bulla and enclosed otic capsule (Refs. 88,109,134). The directional properties of the delphinid receiver are poorly known\* (Refs. 23,88,111). There is some evidence these animals may be able to process separately the signals received on each side (Ref. 19,23,88). Other than its obvious utility for localization, the advantages of this apparent capability are moot.

The external ears of the bats are highly variable in relative size and form (Refs. 70,120,122,131,132). Their shape is often correlated with other behavioral or acoustic features. For example, many small-eared species are fast fliers and emit intense, high

---

\* Measurements performed to date have all employed tone burst signals. Analogous data obtained using an echolocation-like click will probably differ somewhat from these.

frequency sounds. Other species that hunt resting or terrestrial prey emit faint sounds and have relatively large pinnae. An external appendage, the tragus, located along the anterior margin of the external meatus, is prominent in many species (Refs. 70,131,132). Evidence indicates the function(s) of the pinna and tragus is to effect directional sensitivity; the pinna behaves as an attenuator to signals from undesired directions (rather than as an amplifier to sounds from the desired direction), and the tragus increases directional sensitivity in the vertical plane in some unknown fashion (Ref. 64).

Some species move their pinnae during echolocation, some synchronously with pulse emission, and others asynchronously. The purpose of this movement is unknown (see previous section on signal processing). Other species show no apparent ear movements. Directional sensitivity is a complex function of frequency and is dependent upon the shape and orientation of the pinnae. Bilateral symmetry apparently prevails. It seems likely the FM bats exploit the interaction between signal bandwidth and directional sensitivity to localize a target in three-dimensional space (Ref. 64). The CF bats could exploit the FM tail of their signals similarly. It is concluded that frequency/intensity differences between the ears are probably the determinants of sound localization (Refs. 59,97,132). At high signal frequencies directivity is of the order of  $0.75 \text{ dB}/^\circ$  (Ref. 59). Intensity differences of 0.5 to 0.75 dB are discernible (Ref. 57). Thus, it appears that bats should be able to localize objects to within  $1^\circ$  (or less), at least within the conus of radiation.

Owls are characterized by asymmetry in the location of their external auditory meatuses on the head (Ref. 79,107,125); a line connecting their "acoustic centers" makes an angle of  $10^\circ$  to  $15^\circ$  with the horizontal. Directional sensitivity is a complex function

of frequency\* (Refs. 79,107,125). The asymmetry and directional sensitivity interact to permit accurate (and apparently unique) localization of a radiating source in the frontal hemisphere (Refs. 79, 107,125). Upon first detecting a sound in darkness, the owl turns its head toward the source. Once it faces the source, it must hear one additional sound before striking (Ref. 125). Thus, the owl must apparently face the source "directly" to accurately localize it. Whether the owl determines target range before attacking or senses its approach toward the ground along a path of target interception is not known; extant data neither support nor reject either hypothesis (Refs. 79,125).

Synthetic sonars are probably capable of achieving localization accuracies as good as, and perhaps better than, the bats'. However, the relative time required to do so may significantly exceed that required by these animals. Accordingly, further work in this area would appear to be potentially rewarding for sonar technology. The owl's apparent ability to ascertain target range passively is truly exciting, and support of further (and expanded) study of this animal is urged.

---

\* There are minimal data available (Ref. 126) and the validity of some of these data has been questioned (Ref. 79). Also, measurements to date have employed pure tone stimuli. Natural stimuli for owls are probably broadband rustling, crunching, or chewing noises. Measurements performed with these kinds of stimuli would exhibit differences from the pure tone results.

## V. PASSIVE/ACTIVE SONAR UTILIZATION

Some species of bats apparently exploit target radiated noise for early detection and localization of prey (Ref. 62). This is not a species' specific capability, but seems related only to bat/prey size. Absolute hearing sensitivity appears the same for all bats (and, indeed, is equivalent to other mammals). Accordingly, passive sonar capabilities of different species should be comparable in this regard. Large insect prey do radiate sound at intensities adequate to be exploited passively. On the other hand, small insect prey, e.g., fruit flies and mosquitoes, radiate sound at levels too low to be useful at any seemingly reasonable range to effect pursuit action (Ref. 51). The larger species of bats pursue and capture larger insect prey, while the smaller species pursue small insect prey. Therefore, it is likely the larger species rely on target radiated sounds for early detection and localization (there is some experimental evidence supporting this conjecture (Ref. 82)), while it is moot to know to what extent the smaller species do, if they do at all.

Confounding measurement on bats is the phenomenon of spatial or proprioceptive learning observed in the laboratory, but probably occurring to some degree in the wild. Given time to examine and learn their environment acoustically, bats reduce their echolocation (orientation) activity to a minimum and rely heavily upon apparent apperception of their own movements through it (Refs. 1,2,80,93, 104,105). It appears that they may not even attend to the few signals emitted during this activity (Refs. 3,44,80). They may be relying upon passive detection of environmental noises, e.g., the rustling of leaves, or, in the laboratory, detection of reflections of flight generated sounds or of noise generated by scraping or tapping their claws

when crawling along a surface. That is, the bats may be exploiting a passive sonar capability for orientation (navigation) in familiar surroundings.

The owl's reliance upon and ability to use target radiated sounds has been discussed in preceding sections.

It is probable that whales rely (perhaps heavily) on passive detection and localization of prey or other sound radiating objects (e.g., ships), although this hypothesis is a difficult one to test. Available evidence is mostly anecdotal in nature. The killer whale, Orcinus orca, has not been detected echolocating in the wild and must be taught to use its capability in captivity (Ref. 37). The bottle-nose dolphin, T. truncatus, has been observed tracking live fish in its tank passively, and echolocated only when the ambient noise level was raised (the water filtration system was activated) (Refs. 10,37). Accordingly, it can be argued that many animals swimming together in a large herd would generate so much self-noise as to render passive target detection unlikely.

However, observation of a large group of dolphins feeding on a school of fish in relatively turbid water (limit of visibility ~50 cm) led to the conclusion that the dolphins were using only visual and passive acoustic clues to detect and capture individual prey (Ref. 25).

Finally, it has been observed that the dolphin in familiar surroundings performing a routine echolocation task becomes acoustically lazy, emitting apparently only a minimal number of signals requisite for the task, and these at very low intensities (Ref. 10). (A similar phenomenon of habituation to the task has also been observed in bats (Ref. 161).).



## VI. CONCLUSIONS

The preceding overview of biological sonar phenomena is an attempt to place the knowledge of, and potential for, research in this area into perspective from the viewpoint of synthetic sonar technology. The guiding philosophy for this survey was to ascertain and illuminate those areas or activities of biological sonar research which might immediately or eventually influence, alter, or improve synthetic sonar development or design philosophy. This report is brief and, hopefully, to the point. The List of References is comprehensive, but represents only a portion of the literature surveyed. Much of the literature surveyed contributed to an understanding and appreciation of the problems, results, and direction(s) of research on biological sonar systems but was intrinsically of no significance with regard to bionics exploitation.

There is a message evident in bionics research on animal sonars. That is, the ear-brain is, without question, the best available adaptive processor for analyzing acoustic signals. Sonar technology has, for many valid reasons, sought to remove the ear-brain (the listener/operator) from the decision functions (detection/classification) of sonar operation. A baseline of sonar performance utilizing a motivated operator as an acoustic analyzer has never been established. A survey of biological sonars thus begs the recommendation that the utility of the sonar operator be reevaluated and that baseline sonar performance be established using the ear-brain processor.

Five areas of animal sonar research were recommended for continued or additional support because of their apparent potential to provide bionically useful information. Perhaps of most immediate consequence is study of the ability of the owl to localize and

apparently range passively. Investigation of this phenomenon is proceeding with but minimal funding.\* Accordingly, it is recommended that this work receive expanded support.

The second area has been a topic of investigation for many years with varying degrees of activity and with little or no success, primarily due to procedural reasons but often due to instrumental shortcomings. This is the area of signal design for target classification: how are the interrogating signals modified to apparently optimize some target-specific characteristic of the echo? A methodological change was recommended. Given this change, research on both whales and bats appears ready to provide the required data, the work on whales through direct acoustic measurements, that on bats through electrophysiological techniques. Knowledge of the target acoustics is, of course, implicit.

Electrophysiological measurements are a direct approach in attempting to comprehend ear-brain functioning in the analysis of acoustic signals. The potential contribution of work in this area is great, but admittedly long-range. Support in this area, to the extent of maintaining an awareness of activities and results, was recommended. Understanding the animals' ability to localize a target in three-dimensional space is corollary to this work. More is known about bats in this area. Directivity of the external ear structures is a complex function of frequency, apparently permitting unique localization of a target in space with each echo. Knowledge of how bats achieve this localization might be exploited to facilitate analogous measurements by synthetic sonars. Therefore, it was recommended that work in this area be supported, at least to the extent of maintaining an awareness of activities and results.

---

\*The only known research being conducted in this country at this time is by Professor Mark Konishi, Department of Biology, Princeton University, under an NSF grant.

Finally, evolutionary evidence implies the delphinid sonar is optimal for underwater short range, high resolution applications. A bio-analog (electromechanical) simulation has been developed and is being used to examine the potential and limitations of the delphinid signal form (see Fig. 1). However, mechanical generation is inefficient and severely limits versatility. Accordingly, it was recommended that study of the dolphin's signal generator be supported, principally to guide development of possible new means of generating the high intensity, broadband signals required for exploitation of this sonar form, but also as the generator determines the limits of the animal's signal design capabilities.

Appendix II is a tabulation of available sonar and performance data for bats. One hundred three (103) genera and/or species are listed; Table A includes the Old World (Eastern hemisphere) bats; Table B, the New World (Western hemisphere) bats. Entries are alphabetical within each table. There are no obvious consistencies (or inconsistencies, for that matter) or trends evident in these data. What is obvious is that there is a multiplicity of signal forms used to solve many similar and diverse problems, all obviously successfully. What is also apparent is that in many areas there is a dearth of data extant.

Appendix III is a tabulation of available sonar and performance data for whales, dolphins, and porpoises. Nineteen (19) species are listed. Entries are alphabetical; there is no further breakdown. It is immediately apparent from this tabulation that there are appreciably fewer data available for these animals than for bats. The general trend apparent in these data is an inverse correlation between signal frequency (or absolute bandwidth) and physical size of the animal. All species seem to utilize a transient or very short pulse for echolocation.

Bibliographies of the accumulated literature for bats, whales, and other animals are included as Appendices IV, V, and VI, respectively. Entries are alphabetical by (senior) author, and chronological within an author's listing(s). Each entry has been coded using a 6-digit number. The coding key is shown at the beginning of each bibliography.

An annotated bibliography of selected papers, plus a compendium of selected translations of the foreign language literature, will be issued at a later date.

# REFERENCES

1. Airapetiantz, E. Sh., "Bionics Aspects of Research Into Mechanisms Responsible for Spatial Information Echolocation in Bats," *Fiziologicheskii Zh. SSSR Moscow* 54, 368-376 (1967), (Russian; English translation).
2. Ayrapet'yants, E. Sh., A. G. Golubkov, I. V. Yershova, A. R. Zhezherin, V. N. Zvorykin, and V. I. Korolev, "Echolocation Differentiation and Characteristics of Radiated Pulses in Dolphins," *Dokl. Akad. Nauk SSSR* 188, 1197-1199 (1969); JPRS 49479.
3. Airapetiantz, E. Sh., and A. I. Konstantinov, "On the Interaction of Analyzers in the Echolocation Activities of Bats," in the collection *Issledovaniye Apparator Signalizatsii Mozga* (Nauka Publishing House, Leningrad, 1967), (Russian; English translation).
4. Ayrapet'yants, E. Sh., and A. I. Konstantinov, Echolocation in Nature (Nauka Publishing House, Leningrad, 1970), Ch. 17 and Conclusion; JPRS 51511.
5. Airapetiantz, E. Sh., and A. I. Konstantinov, "Physiological Investigations of Ultrasonic Echolocation in Animals," paper presented at the 25th International Congress of Physiol. Sciences, Munich (1971).
6. Airapetianz, E. Sh., A. I. Konstantinov, and D. P. Matjushkin, "Brain Echolocation Mechanisms and Bionics," *Acta Physiol. Acad. Sci. Hung.* 35, 1-17 (1969).
7. Altes, R. A., and E. L. Titlebaum, "Bat Signals as Optimally Doppler Tolerant Waveforms," *J. Acoust. Soc. Amer.* 48, 1014-1017 (1970).
8. Altes, R. A., "Computer Derivation of Some Dolphin Echolocation Signals," *Science* 173, 912-914 (1971).
9. Andersen, S., "Auditory Sensitivity of the Harbor Porpoise Phocoena phocoena," in Investigations on Cetacea, G. Pilleri (ed.), (Benteli AG, Berne, Switzerland, 1970), Vol. II, pp. 256-259.
10. Applied Research Laboratories, The University of Texas at Austin, and Naval Undersea Center, San Diego, unpublished data (1969).
11. Applied Research Laboratories, The University of Texas at Austin, and Naval Undersea Center, San Diego, unpublished data (1972).
12. Applied Research Laboratories, The University of Texas at Austin, Quarterly Progress Report No. 3 under Contract N60530-68-C-0882, for the period 1 December 1968 through 28 February 1969.

13. Applied Research Laboratories, The University of Texas at Austin, Quarterly Progress Reports No. 2 and No. 3 under Contract N66001-70-C-0268, for the period 16 December 1969 through 15 June 1970.
14. Applied Research Laboratories, The University of Texas at Austin, Quarterly Progress Reports under Contract N00123-71-C-1593, for the period 3 May 1971 through 1 May 1972.
15. Backus, R. H., and W. E. Schevill, "Physeter Clicks," in Whales, Dolphins, and Porpoises, K. S. Norris, (ed.) (University of California Press, Berkeley, 1966), pp. 510-528.
16. Bagdonas, A., V. M. Bel'kovich, and N. L. Krushinskaya, "Interaction of Analyzers in Dolphins During Discrimination of Geometrical Figures Under Water," J. Higher Neural Activity 20, 1070-1075 (1970), (Russian; English translation).
17. Barta, R. E., "Acoustical Pattern Discrimination by an Atlantic Bottlenose Dolphin," Naval Undersea Center, San Diego, California, (unpublished manuscript), (1969).
18. Batteau, D. W., "Theories of Sonar Systems and Their Application to Biological Organisms," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 2, pp. 1033-1091.
19. Bel'kovich, V. M., "Anatomy and Function of the Ear in Dolphins," Zoologicheskii Zh. 2, 275-282 (1970); JPRS 50253.
20. Bloedel, P., "Observations on the Life Histories of Panama Bats," J. Mamm. 36, 232-235 (1955).
21. Bradbury, J. W., "Target Discrimination by the Echolocating Bat Vampyrum spectrum," J. Exp. Zool. 173, 23-46 (1970).
22. Bradbury, J. W., and F. Nottebohm, "Use of Vision by the Little Brown Bat, Myotis lucifugus, under Controlled Conditions," Anim. Behav. 17, 480-485 (1969).
23. Bullock, T. H., A. D. Grinnell, E. Ikezono, K. Kameda, Y. Katsuki, M. Nomoto, O. Sato, N. Suga, and K. Yanagisawa, "Electrophysiological Studies of Central Auditory Mechanisms in Cetaceans," Z. vergl. Physiol. 52, 117-156 (1968).
24. Bullock, T. H., and S. H. Ridgway, "Evoked Potentials in the Central Auditory System of Alert Porpoises to Their Own and Artificial Sounds," J. Neurobiol. 3, 79-99 (1972).
25. Busnel, R. G., personal communication (1972).

26. Busnel, R. G., A. Dziedzic, and S. Andersen, "Seuils de Perception du Systeme Sonar du Marsouin Phocaena phocaena, in Function du Diametre d'un Obstacle Filiforme," C. R. Acad. Sci. 260, 295-297 (1965), (in French).
27. Busnel, R. G., and A. Dziedzic, "Resultants Metrologiques Experimentaux de l'Echolocation chez le Phocaena phocaena, et leur Comparaison avec ceux de Certaines Chauves-Sousis," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 307-338 (in French).
28. Cahlander, D. A., J. J. G. McCue, and F. A. Webster, "The Determination of Distance by Echolocating Bats," Nature 201, 544-546 (1964).
29. Chase, J., and R. A. Suthers, "Visual Obstacle Avoidance by Echolocating Bats," Anim. Behav. 17, 201-207 (1969).
30. Dalland, J. I., "Hearing Sensitivity in Bats," Science 150, 1185-1186 (1965).
31. Diercks, K. J., F. L. Weisser, and W. E. Evans, "Analysis of Short Pulse Echoes from Copper Plates," J. Acoust. Soc. Amer. 42, 1211(A) (1967).
32. Diercks, K. J., W. W. Ryan, E. E. Mikeska, and F. L. Weisser, "Listener Discrimination of Broadband FM Echoes from Simple Geometric Targets," Defense Research Laboratory Technical Memorandum No. 68-22 (DRL-TM-68-22), Defense Research Laboratory, The University of Texas at Austin, (25 November 1968), 50 pages.
33. Diercks, K. J., R. T. Trochta, C. F. Greenlaw, III, and W. E. Evans, "Recording and Analysis of Dolphin Echolocation Signals," J. Acoust. Soc. Amer. 49, 1729-1732 (1971).
34. Diercks, K. J., and R. T. Trochta, "Animal Sonar: Measurements and Meaning," J. Acoust. Soc. Amer. 51, 133(A) (1972) (to be published in J. Acoust. Soc. Amer.).
35. Dubrovsky, N. A., A. A. Titov, P. S. Krasnov, V. P. Babkin, V. M. Lekontsev, and G. V. Nikolenko, "Investigation of the Permission (sic) Capacity of the Black Sea Tursiops truncatus Echolocation Apparatus," Trudy Akust. Inst. 10, 163-181 (1970), (Russian; discussed in Ref. 153).
36. Escudie, B., A. Hellion, and A. Dziedzic, "Quelques Resultats dans l'etude des Sonars Biologiques Ariens et Marins par Traitement du Signal et Analyse Spectrale," in Proc. Troisieme Colloque sur le Traitement du Signal et ses Applications, Nice, France, 1-5 June 1971, 533-557.

37. Evans, W. E., personal communication (1972).
38. Evans, W. E., and B. A. Powell, "Discrimination of Different Metallic Plates by an Echolocating Delphinid," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 363-382.
39. Evans, W. E., and J. H. Prescott, "Observations of the Sound Production Capabilities of the Bottlenose Porpoise; a Study of Whistles and Clicks," Zoologica 47, 121-128 (1962).
40. Evans, W. E., W. W. Sutherland, and R. G. Beil, "Directional Characteristics of Delphinid Sounds," in Marine Bio-Acoustics, W. N. Tavolga (ed.), (The Macmillan Company, New York, 1964), pp. 353-372.
41. Evans, W. E., and E. C. Evans, III, "Echolocation of Aquatic Mammals Based on Experimental Evidence from Marine and Freshwater Cetaceans," Naval Undersea Center, San Diego, California, unpublished manuscript (1971).
42. Griffin, D. R., "Bat Sounds Under Natural Conditions with Evidence for Echolocation of Insect Prey," J. Exp. Zool. 123, 435-466 (1953).
43. Griffin, D. R., "Acoustic Orientation in the Oilbird, Steatornis," Proc. Nat. Acad. Sci. 39, 884-893 (1953).
44. Griffin, D. R., Listening in the Dark (Yale University Press, New Haven, 1958).
45. Griffin, D. R., "Comparative Studies of the Orientation Sounds of Bats," Symp. Zool. Soc. London 7, 61-72 (1962).
46. Griffin, D. R., "Discriminative Echolocation by Bats," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 273-305.
47. Griffin, D. R., personal communication (1972).
48. Griffin, D. R., and A. Novick, "Orientation of Neotropical Bats," J. Exp. Zool. 130, 251-300 (1955).
49. Griffin, D. R., A. Novick, and M. Kornfield, "The Sensitivity of Echolocation in the Fruit Bat, Rousettus," Biol. Bull. 115, 107-113 (1958).
50. Griffin, D. R., and A. D. Grinnell, "Ability of Bats To Discriminate Echoes from Louder Noises," Science 128, 145-146 (1958).



51. Griffin, D. R., F. A. Webster, and C. R. Michael, "The Echolocation of Flying Insects by Bats," *Anim. Behav.* 8, 141-154 (1960).
52. Griffin, D. R., D. C. Dunning, D. A. Cahlander, F. A. Webster, J. D. Pye, M. Flinn, and A. Pye, "Correlated Orientation Sounds and Ear Movements of Horseshoe Bats," *Nature* 196, 1185-1188 (1962).
53. Griffin, D. R., J. J. G. McCue, and A. D. Grinnell, "The Resistance of Bats to Jamming," *J. Exp. Zool.* 152, 229-250 (1963).
54. Griffin, D. R., J. H. Friend, and F. A. Webster, "Target Discrimination by the Echolocation of Bats," *J. Exp. Zool.* 158, 155-168 (1965).
55. Griffin, D. R., and R. A. Suthers, "Sensitivity of Echolocation in Cave Swiftlets," *Biol. Bull.* 139, 495-501 (1970).
56. Grimley, W. K., and E. J. Risness, in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 304-306.
57. Grinnell, A. D., "The Neurophysiology of Audition in Bats: Intensity and Frequency Parameters," *J. Physiol.* 167, 38-66 (1963).
58. Grinnell, A. D., "The Neurophysiology of Audition in Bats: Temporal Parameters," *J. Physiol.* 167, 67-96 (1963).
59. Grinnell, A. D., "The Neurophysiology of Audition in Bats: Directional Localization and Binaural Interaction," *J. Physiol.* 167, 97-113 (1963).
60. Grinnell, A. D., "Mechanisms of Overcoming Interference in Echolocating Animals," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 451-478.
61. Grinnell, A. D., "Comparative Auditory Neurophysiology of Neotropical Bats Employing Different Echolocation Signals," *Z. vergl. Physiol.* 68, 117-153 (1970).
62. Grinnell, A. D., personal communication (1972).
63. Grinnell, A. D., and D. R. Griffin, "The Sensitivity of Echolocation in Bats," *Biol. Bull.* 114, 10-22 (1958).
64. Grinnell, A. D., and V. S. Grinnell, "Neural Correlates of Vertical Localization by Echolocating Bats," *J. Physiol.* 181, 830-851 (1965).
65. Grinnell, A. D., and S. Hagiwara, "Adaptations of the Auditory Nervous System for Echolocation," *Z. vergl. Physiol.* 76, 41-81 (1972).

66. Grummon, R. A. and A. Novick, "Obstacle Avoidance in the Bat, Macrotus Mexicanus," Physiol. Zool. 36, 361-369 (1963).
67. Gurevich, V. S., "Echolocation Discrimination of Geometric Figures in the Dolphin, Delphinus delphis," Moscow, Vestnik Moskovskogo Universiteta, Biologiya, Pochovedeniye 3, 109-112 (1969); JPRS 49281.
68. Hall, J. D. and C. S. Johnson, "Auditory Thresholds of a Killer Whale, Orcinus orca Linnaeus," J. Acoust. Soc. Amer. 51, 515-517 (1972).
69. Henson, O'D. W., Jr., "The Perception and Analysis of Biosonar Signals by Bats," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 2, pp. 949-1003.
70. Henson, O'D. W. Jr., "The Ear and Audition," in The Biology of Bats, W. A. Wimsatt (ed.), (Academic Press, New York, 1970), pp. 181-263.
71. Henson, O'D. W., Jr. and G. D. Pollak, "A Technique for Chronic Implantation of Electrodes in the Cochleae of Bats," unpublished manuscript (1972).
72. Jacobs, D. W. and J. D. Hall, "Auditory Thresholds of a Fresh Water Dolphin, Inia geoffrensis Blainville," J. Acoust. Soc. Amer. 51, 530-533 (1972).
73. Johnson, C. S., "Sound Detection Thresholds in Marine Mammals," in Marine Bio-Acoustics, W. N. Tavolga (ed.), (The Macmillan Company, New York, 1967), Vol. 2, pp. 247-260.
74. Johnson, C. S., in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 384-398.
75. Johnson, C. S., "Relation Between Absolute Threshold and Duration of Tone Pulses in the Bottlenosed Porpoise," J. Acoust. Soc. Amer. 43, 757-763 (1968).
76. Johnson, C. S., "Masked Tonal Thresholds in the Bottlenosed Porpoise," J. Acoust. Soc. Amer. 44, 965-967 (1968).
77. Kellogg, W. N., "Echo Ranging in the Porpoise," Science 128, 981-988 (1958).
78. Kellogg, W. N., "Size Discrimination by Reflected Sound in a Bottlenosed Porpoise," J. Acoust. Soc. Amer. 52, 509-514 (1959).
79. Konishi, M., personal communication (1972).

80. Konstantinov, A. I., "Principles of Ultrasound Spatial Orientation in Bats," *Vopr. Sravnit. Fiziol. Analiz. Leningrad* 2, 93-111 (1966), (Russian; English translation).
81. Konstantinov, A. I., "Relationship between Auditory Perception and Echolocation During Hunting in *Myotis oxynathus*," *Zhurn. Evolyut. Fiziol. i Biokhim.* 5, 566-572 (1969), (Russian; English translation).
82. Konstantinov, A. I., and N. I. Akhmarova, "Target Discrimination by Echolocation of Bats," *Nau. Dok. Vys. Shk. Biol. Nauki* 11, 22-28 (1968), (Russian; English translation).
83. Konstantinov, A. I., N. F. Mel'kinov, and A. A. Titov, "On the Abilities of Dolphins To Recognize Objects," *Tez. Dokl. II Respubl. Konf. po Bionike, Kiev*, 57-59 (1968), (Russian; English translation).
84. Konstantinov, A. I., B. V. Sokolov and I. M. Stosman, "Comparative Research in Echolocation Sensitivity in Bats," *Dan SSSR* 175, 1418-1421 (1967), (Russian; English translation).
85. Konstantinov, A. I., and B. V. Sokolov, "Characteristics of Ultrasonic Orientation Signals in Horseshoe Bats (*Rhinolophidae*)," *Zh. Evol. Biok. i Fiziol.* 5, 90-97 (1969), (Russian; English translation).
86. Konstantinov, A. I., N. N. Sanotskaya, and N. N. Sokolova, "Frequency-Threshold Characteristics of the Auditory System of Bats Measured by the Conditioned Reflex Method," *J. Higher Nervous Act.* 21, 535-541 (1971).
87. Kulzer, E., "Flying Fox Produces Orientation Sounds with Tongue," *Naturwiss.* 43, 117-118 (1956).
88. McCormick, J. G., E. G. Wever, J. Palin, and S. H. Ridgway, "Sound Conduction in the Dolphin Ear," *J. Acoust. Soc. Amer.* 48, 1418-1428 (1970).
89. McCue, J. J. G., "Aural Pulse Compression by Bats and Humans," *J. Acoust. Soc. Amer.* 40, 545-548 (1966).
90. McCue, J. J. G., "Signal Processing by the Bat," *J. Aud. Res.* 9, 100-107 (1969).
91. Mogus, M. A., "Theories of Bat Echolocation," Ordnance Research Laboratory Technical Memorandum No. 657.2341-02, Ordnance Research Laboratory, Pennsylvania State University (February 1967), 62 pages. AD 650476.
92. Mogus, M. A., "A Theoretical Approach to Bat Echolocation," Ph.D. Dissertation, Pennsylvania State University (1970), University Microfilms, Inc., Ann Arbor, Michigan.

93. Mohres, F. P., "On the Orientation of Bats," *Natur und Volk* 80, 153-161 (1950), (German; English translation).
94. Mohres, F. P., "Concerning a New Type of Ultrasonic Orientation in Bats," *Verh. Dtsch. Zool.* 179-186 (1951), (German; English translation).
95. Mohres, F. P., "The Ultrasonic Orientation of Bats," *Naturwiss.* 39, 273-279 (1952), (German; English translation).
96. Mohres, F. P., "Ultrasonic Orientation in Flying Fox Bats," *Naturwiss.* 40, 536-537 (1953), (German; English translation).
97. Mohres, F. P., "About the Ultrasonic Orientation of the Horseshoe Bat (*Chiroptera-Rhinolophinae*)," *Z. vergl. Physiol.* 34, 547-588 (1953), (German; English synopsis).
98. Mohres, F. P., "Ultrasonic Orientation in Megadermatid Bats," in *Animal Sonar Systems: Biology and Bionics*, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 115-126.
99. Mohres, F. P., "General Characters of Acoustic Orientation Signals and Performance of Sonar in the Order Chiroptera," in *Animal Sonar Systems: Biology and Bionics*, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 401-407.
100. Mohres, F. P., and G. Neuweiler, "Die Ultraschallorientierung der Grossblatt-Fledermäuse (*Chiroptera-Megadermatidae*)," *Z. vergl. Physiol.* 53, 195-227 (1966), (German; English synopsis).
101. Morozov, V. P., A. I. Akopian, V. I. Burdin, K. A. Zaytseva, and Yu. A. Sokovykh, "Repetition Rate of Ranging Signals of Dolphins as a Function of Distance to Target," *Biofizika* 17, 139-144 (1972); JPRS 55729.
102. Neuweiler, G., "Interaction of Other Sensory Systems with the Sonar System," in *Animal Sonar Systems: Biology and Bionics*, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 509-529.
103. Neuweiler, G., "Neurophysiological Investigations in the Echo-orientation of the Greater Horseshoe Bat *Rhinolophus ferrum equinum*," *Z. vergl. Physiol.* 67, 273-306 (1970), (German; English translation).
104. Neuweiler, G., and F. P. Mohres, "Die Rolle des Ortsgedächtnisses bei der Orientierung der Grossblatt-Fledermäuse *Megaderma lyra*," *Z. vergl. Physiol.* 57, 147-171 (1967), (German; English synopsis).

105. Neuweiler, G., and F. P. Mohres, "Role of Spacial (sic) Memory in the Orientation," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 129-139.
106. Neuweiler, G., G. Schuller, and H. V. Schnitzler, "On- and Off-Responses in the Inferior Colliculus of the Greater Horseshoe Bat to Pure Tones," Z. vergl. Physiol. 74, 57-63 (1971).
107. Norberg, A., "Physical Factors in Directional Hearing in Aegolius funereus (Linne) (Strigiformes), with Special Reference to the Significance of the Asymmetry of the External Ears," Arkiv for Zoologi 20, 181-204 (1968).
108. Norris, K. S., "Some Problems of Echolocation in Cetaceans," in Marine Bio-Acoustics, W. N. Tavolga (ed.), (The Macmillan Company, New York, 1964), pp. 317-336.
109. Norris, K. S., "Echolocation of Marine Mammals," in Biology of Marine Mammals, H. T. Andersen (ed.), (Academic Press, New York,
110. Norris, K. S., personal communication (1972).
111. Norris, K. S., J. H. Prescott, P. V. Asa-Dorian, and P. Perkins, "Experimental Demonstration of Echolocation Behavior in the Porpoise Tursiops truncatus (Montagu)," Biol. Bull. 20, 163-176 (1961).
112. Norris, K. S., and W. E. Evans, "Directionality of Echolocation in the Rough-Tooth Porpoise Steno bredanensis (Lesson)," in Marine Bio-Acoustics, W. N. Tavolga (ed.), (The Macmillan Company, New York, 1967), Vol. 2, pp. 305-316.
113. Norris, K. S., W. E. Evans, and R. N. Turner, "Echolocation in an Atlantic Bottlenose Porpoise During Discrimination," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, 409-437.
114. Norris, K. S., K. J. Dormer, J. Pegg, and G. J. Liese, "The Mechanism of Sound Production and Air Recycling in Porpoises: A Preliminary Report," unpublished manuscript. The Oceanic Institute, Waimanalo, Hawaii (1972), 13 pages.
115. Novick, A., "Acoustic Orientation in the Cave Swiftlet," Biol. Bull. 117, 497-503 (1959).
116. Novick, A., "Orientation of Paleotropical Bats: I. Microchiroptera," J. Exp. Zool. 138, 81-154 (1958).

117. Novick, A., "Pulse Duration in the Echolocation of Insects by the Bat Pteronotus," Ergeb. der Biol. 26, 21-26 (1963).
118. Novick, A., "Orientation in Neotropical Bats: II. Phyllostomatidae and Desmodontidae," J. Mamm. 44, 44-56 (1963).
119. Novick, A., "Echolocation of Flying Insects by the Bat Chilonycteris psilotis," Biol. Bull. 128, 297-314 (1965).
120. Novick, A., "Echolocation in Bats: Some Aspects of Pulse Design," Am. Sci. 59, 198-209 (1971).
121. Novick, A., and N. Leen, The World of Bats (Holt, Reinhart, and Winston, New York, 1970).
122. Novick, A., personal communication (1972).
123. Novick, A., and D. R. Griffin, "Laryngeal Mechanisms in Bats for the Production of Orientation Sounds," J. Exp. Zool. 148, 125-145 (1961).
124. Novick, A., and J. R. Vaisnys, "Echolocation of Flying Insects by the Bat Chilonycteris parnelli," Biol. Bull. 127, 478-488 (1964).
125. Payne, R., "Acoustic Location of Prey by Barn Owls (Tyto alba)," J. Exp. Biol. 54, 535-573 (1971).
126. Peff, T. C., and J. A. Simmons, "Horizontal Angle Resolution by Echolocating Bats," J. Acoust. Soc. Amer. 51, 2063-2065 (1972).
127. Pollak, G., O. W. Henson, Jr., and A. Novick, "Cochlear Microphonic Audiograms in the 'Pure Tone' Bat Chilonycteris parnellii parnellii," Science 176, 66-68 (1972).
128. Penner, R. H., and A. E. Murchison, "Experimentally Demonstrated Echolocation in the Amazon River Porpoise Inia geoffrensis (Blainville)," Naval Undersea Center Tech. Publ. No. 187, June 1970, 28 pages.
129. Purves, P. E., "Anatomical and Experimental Observations on the Cetacean Sonar System," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 197-270.
130. Pye, A., "Structure of Cochlea in Chiroptera III. Microchiroptera: Phyllostomatoidea," J. Morph. 121, 241-254 (1967).
131. Pye, J. D., "Echolocation by Bats," Endeavour 20, 101-111 (1961).

132. Pye, J. D., "Hearing in Bats," in Symp. CIBA Found. "Hearing Mechanisms in Vertebrates," de Reuck and Knight (eds.), (1968), pp. 66-88.
133. Pye, J. D., and L. H. Roberts, "Ear Movements in a Hipposiderid Bat," Nature 225, 285-286 (1970).
134. Reysenbach De Haan, F. W., "Listening Underwater: Thoughts on Sound and Cetacean Hearing," in Whales, Dolphins, and Porpoises, K. S. Norris (ed.), (University of California Press, Berkeley, 1966).
135. Reznik, A. M., V. M. Skorniyakov, and A. G. Chupakov, "Location Activity of Black Sea Tursiops truncatus Being Presented Targets," Trudy Akusticheskogo Inst. 12, 116-120 (1970), (Russian; discussed in Ref. 153).
136. Roeder, K. D., "Echoes of Ultrasonic Pulses from Flying Moths," Biol. Bull. 124, 200-209 (1963).
137. Romanenko, E. V., "Underwater Echolocation (SONAR) Capacity of Dolphins (Review)," Sov. Phys. Acoust. 10, 331-342 (1965).
138. Romanenko, E. V., A. G. Tomilin, and B. A. Artemenko, "On the Question of Sound Formation and the Directing of Sounds in Dolphins," in the collection Bionika, 369-373 (1965), (Russian; English translation).
139. Schevill, W. E., and W. A. Watkins, "Sound Structure and Directionality in Orcinus (Killer Whale)," Zoologica 51, 71-76 (1966).
140. Schnitzler, H. U., "Discrimination of Thin Wires by Flying Horseshoe Bats (Rhinolophidae)," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.), (France, 1967), Vol. 1, pp. 69-87.
141. Schnitzler, H. U., "Die Ultraschall-Ortungslaute der Hufeisen Fledermause (Chiroptera-Rhinolophidae) in Verschiedenen Orientierungssituationen," Z. vergl. Physiol. 57, 376-408 (1968), (German; English summary).
142. Schnitzler, H. U., "Echoortung bei der Fledermaus Chilonycteris rubiginosa," Z. vergl. Physiol. 68, 25-38 (1970), (German; English translation).
143. Schnitzler, H. U., "Comparison of the Echolocation Behaviors in Rhinolophus ferrum equinum and Chilonycteris rubiginosa," Bijdragen Tot de Dierkunde 40, 77-79 (1970).
144. Schnitzler, H. U., G. Schuller, and G. Neuweiler, "Antworten der Colliculus inferior der Fledermaus Rhinolophus euryale auf Tonale Reizung," Die Naturwiss. 12, 627.

145. Simkin, G. N., and N. D. Patlyakevich, "Motion of Bats Toward Plane Targets and the Nature of Changes of Echolocating Signals in the Process of Target Selection and Recognition," *Biologicheskiye Nauki* 3, 48-57 (1969); JPRS 48123.
146. Simkin, G. N., "Echolocation Process in Oxyrathous Bats in Free-Flight," *Biologicheskiye Nauki* 3, 61-71 (1970); JPRS 50665.
147. Simmons, J. A., "Depth Perception by Sonar in the Bat Eptesicus fuscus," Ph.D. Dissertation, Princeton University (December 1969); University Microfilms, Inc., Ann Arbor, Michigan.
148. Simmons, J. A., "Acoustic Radiation Patterns for the Echolocating Bats Chilonycteris rubiginosa and Eptesicus fuscus," *J. Acoust. Soc. Amer.* 46, 1054-1056 (1969).
149. Simmons, J. A., "Echolocation in Bats: Signal Processing of Echoes for Target Range," *Science* 171, 925-928 (1971).
150. Simmons, J. A., personal communication (1972).
151. Simmons, J. A., "Response of the Doppler Sonar System in the Bat Rhinolophus ferrum equinum," unpublished manuscript (1972), (submitted for publication in *J. Acoust. Soc. Amer.*).
152. Simmons, J. A., and J. A. Vernon, "Echolocation: Discrimination of Targets by the Bat Eptesicus fuscus," *J. Exp. Zool.* 176, 315-328 (1971).
153. Sokolov, V., "Cetacean Research in the USSR," in Investigations on Cetacea, G. Pilleri (ed.) (Benteli AG, Berne, Switzerland, 1971), Vol. III(2), pp. 317-346.
154. Sokolov, B. V., and A. K. Makarov, "The Direction of Ultrasonic Orientational Radiation of the Large Rhinolophidae and the Role of Nasal Outgrowths in its Formation," *Sci. Rpts. of the Higher Sch., Biol. Sci.* 7, 37-44 (1971), (Russian; English translation).
155. Strother, G. K., and M. A. Mogus, "Acoustical Beam Patterns for Bats: Some Theoretical Considerations," *J. Acoust. Soc. Amer.* 48, 1430-1432 (1970).
156. Supin, A. Ya., and M. N. Sukchoruchko, "The Determination of Auditory Thresholds in Phocoena phocoena by the Method of Skin-Galvanic Reaction," *Trudy Akousticheskogo Inst.* 12, 194-199 (1970), (Russian; discussed in Ref. 153).



157. Suthers, R. A., "Acoustic Orientation by Fish-Catching Bats," J. Exp. Zool. 158, 319-348 (1965).
158. Suthers, R. A., "Comparative Echolocation by Fishing Bats," J. Mamm. 48, 79-87 (1967).
159. Suthers, R. A., J. Chase, and B. Braford, "Visual Form Discrimination by Echolocating Bats," Biol. Bull. 137, 535-546 (1969).
160. Van Bergeijk, W. A., "Sonic Pulse-Compression in Bats and People: A Comment," J. Acoust. Soc. Amer. 36, 594-597 (1964).
161. Vernon, J., and E. A. Peterson, "Echolocation Signals in the Free-Tailed Bat, Tadarida mexicana," J. Aud. Res. 5, 317-330 (1965).
162. Vernon, J., and E. A. Peterson, "Hearing in the Vampire Bat, Desmodus rotundus murinus, as shown by Cochlear Potentials," J. Aud. Res. 6, 181-187 (1966).
163. Walker, E. P., Mammals of the World (The Johns Hopkins Press, Baltimore, 1968), 2nd Ed., Vol. 1, pp. 182-392; Vol. 2, pp. 1083-1145.
164. Webster, F. A., "Active Energy Radiating Systems: The Bat and Ultrasonic Principles II; Acoustical Control of Airborne Interceptions by Bats," in Proc. of the Int'l. Cong. on Tech. and Blindness, L. L. Clark (ed.), (Amer. Found. for Blind, New York, 1963), pp. 49-135.
165. Webster, F. A., in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.) (France, 1967), Vol. 1, pp. 626-670.
166. Webster, F. A., "Interception Performance of Echolocating Bats in the Presence of Interference," in Animal Sonar Systems: Biology and Bionics, R. G. Busnel (ed.) (France, 1967), Vol. 1, pp. 673-713.
167. Webster, F. A., and O. G. Brazier, "Experimental Studies on Target Detection, Evaluation and Interception by Echolocating Bats," AMRL Tech. Rpt. 65-172, (November 1965), 144 pages. AD 628055.
168. Webster, F. A., and O. G. Brazier, "Experimental Studies on Echolocation Mechanisms in Bats," AMRL Tech. Rpt. 67-192, May 1968, 165 pages. AD 673373.
169. Webster, F. A., and O. G. Brazier, "Echolocation Investigations on Bats and Humans: Target Localization and Evaluation," AMRL Tech. Rpt. 68-155, September 1969, 73 pages. AD 697070.

170. Evans, W. E., "A Discussion of Echolocation by Cetaceans Based on Experiments with Marine Delphinids and One Species of Fresh-Water Dolphin," Naval Underwater Center, San Diego, unpublished manuscript (to be published in J. Acoust. Soc. Amer.).
171. Jacobs, D. W., "Auditory Frequency Discrimination in the Atlantic Bottlenose Dolphin, Tursiops truncatus Montagu: A Preliminary Report," J. Acoust. Soc. Amer. 52, 696-698 (1972).
172. Pye, J. D., "Bimodal Distribution of Constant Frequencies in Some Hipposiderid Bats (Mammalia: Hipposideridae)," J. Zool. Lond. 166, 323-335 (1972).
173. Roberts, L. H., "Variable Resonance in Constant Frequency Bats," J. Zool. Lond. 166, 337-348 (1972).
174. Simmons, J. A., "The Sonar Receiver of the Bat," Ann. N.Y. Acad. Sci. 188, 161-174 (1971).
175. Tomilin, A. G., Istoriya Slepogo Kashalota (History of the Blind Sperm Whale), Izd. Nauka, Moscow (1965); Rough Draft Translation prepared by Translation Div., Foreign Technology Div., WP-AFB, Ohio, FTD-HT-66-434, (5 December 1966) pp. 208.

**APPENDIX I**

**BIOSONAR MEMBERSHIP CONTACTED**

## APPENDIX I

### Biosonar Membership Contacted

Professor K. S. Norris, The University of California at Los Angeles.

Professor Norris is presently attempting to establish a facility for Cetacean research, including echolocation, at The University of California Santa Cruz campus.

Professor A. D. Grinnell, The University of California at

Los Angeles. Professor Grinnell has been active in electrophysiological investigations of neural correlates of echolocation phenomena in bats. One of his students, Miss Pat Brown, is examining the acoustic signals employed by bats for communication.

Professor A. Novick, Dr. O'D. W. Henson, Dr. George Pollak,

Yale University. Professor Novick is completing a chapter on echolocation by bats to appear in Vol. 3 of The Biology of Bats, W. A. Wimsatt (ed.). Drs. Henson and Pollak are using chronic implants of electrodes to measure electrophysiological correlates of echolocation in free-flying bats.

Professor M. Konishi, Princeton University. Professor Konishi is measuring the ability of owls to localize and range upon target radiated noises, under various experimental conditions. He is the only person in the United States known to be studying target localization by owls. From the viewpoint of sonar technology, his results to date appear especially significant.

**Preceding page blank**

Dr. James G. McCormick, Bowman-Gray School of Medicine, Wake-Forest University. Dr. McCormick is a pioneer in electrophysiological measurements of hearing phenomena in dolphins. He is presently examining the effects of hyperbarism on hearing in mammals.

Professor J. A. Simmons, Washington University. Professor Simmons is using artificial echoes to measure abilities of bats to resolve target range differences and to compensate for Doppler shifts caused by target and/or platform motion. He is attempting to instrument a fast response computer system to replay artificial echoes to echolocating bats at a programmed S/N value to examine the bat's ability to overcome jamming and the signal and/or noise parameters most affecting it. He is presently studying the ability of bats to resolve target aspect. A post-doctoral student, Dr. Glenis Long, working in Griffin's laboratory at The Rockefeller University, but primarily under Simmons' guidance, is measuring the ability of bats to resolve spectral differences in artificial echoes.

Professor N. Suga, Washington University. Professor Suga has been and continues to be active in electrophysiological measurements of both sound projecting and receiving phenomena in bats. He has recently demonstrated the existence of neural attenuation in the auditory brain during sound transmission, analogous to blanking in synthetic sonars, and has been able to electrically stimulate the bat brain to cause sound generation, a phenomenon being exploited to examine Doppler compensation by the CF bats.

Professor D. R. Griffin, The Rockefeller University. Professor Griffin is not presently actively involved in echolocation research. He is providing guidance for two post-doctoral students working in his laboratory: Dr. Glenis Long, mentioned above with Simmons, and Dr. Peter Hollander, who is attempting to measure the space-frequency characteristics of the bat's sound field. A third post-doctoral student, Dr. E. R. Buckler, from the University of Montana, will be in Professor Griffin's laboratory during the 1972-1973 academic year to study insect catching behavior of bats.

Dr. R. G. Busnel, Director, Laboratory of Physiological Acoustics, Jouy-en-Josas, France, and American Museum of Natural History. The French laboratory does research on all aspects of animal sound generation, reception, and correlated behavior, only a small part of which is related to classical echolocation. They were quite active at one time in recording and analysis of dolphin echolocation signals and behavior, but their efforts have been severely curtailed recently by lack of funds. The team they have applied to the problem consists of: Dr. J. C. Levy, Mathematician, currently developing a model for echolocation; Dr. A. Dziedzic, Engineer, responsible for the development and implementation of instrumentation; Dr. B. Escudie, Engineer, a student of Mermoz, and an enthusiastic advocate of correlational processing; and Dr. Busnel, Biologist. The team appears well-rounded and quite capable, but for reasons stated, has drifted out of the mainstream of echolocation research on dolphins. Dr. Levy is at the Jouy laboratory. Drs. Escudie and Dziedzic are at the Institute de Chemie et Physique Industrielle, Lyon. Dr. Escudie is presently working closely with Professor Simmons in analyses of bat signals.

Professor R. A. Suthers, Indiana University. Professor Suthers' interests have, in recent years, turned to problems of determining visual capabilities and usage in bats. He has begun investigation of the interaction between vision and echolocation as manifested by electrophysiological measurements performed in the inferior colliculus of the brain. Professor Suthers has one graduate student (name and academic level unknown) measuring the minimum audible angle (MAA) discrimination ability of bats. Professor Suthers, with Professor Suga, is writing a review of bat echolocation to appear in book form (Academic Press) late this year or early next year.

Dr. Donna J. Howell, Auditory Research Laboratories, Princeton University. Dr. Howell is beginning a post-doctoral fellowship at ARL (Princeton), continuing Simmons' work on target ranging by bats.

Miss Donna McDonald, University of Hawaii. Miss McDonald is attempting to measure MAA discriminability of the bottlenose dolphin. Her work is being performed at the NUC/Hawaii facility under supervision of Dr. W. W. L. Au.

Dr. E. C. Evans, III, Dr. W. W. L. Au, Mr. B. A. Powell, Mr. R. H. Penner, Mr. E. A. Murchison, NUC/Hawaii Laboratory.

Dr. C. S. Johnson, Mr. W. E. Evans, NUC/San Diego, California.

#### Foreign Membership Contacted:

Professor J. D. Pye, Kings College, University of London.

Professor Pye for the past few years has been endeavoring to record and catalog the echolocation sounds of as many different species of bats as possible in the wild. To date, he has

catalogued some 100 species. He has a doctoral student, L. H. Roberts, investigating the mechanics of signal generation and control by bats.

Dr. H. U. Schnitzler and Professor G. Neuweiler, University of Tübingen, Germany (at University of Frankfurt, beginning 1 January 1973). The efforts of this group are directed toward a behavioral and electrophysiological understanding of echolocation by the bat Rhinolophus ferrum-equinum. Immediately, an attempt will be made to explain the neurophysiological mechanism of Doppler compensation by this species of bat. Two students of Dr. Schnitzler's, E. Flieger and G. Schuller, are examining the echolocation abilities of this species of bat, and its apparent use of selected portions of its signal for location and ranging.

Professor S. Andersen, Odense University, Denmark. Professor Andersen has made both behavioral and acoustic measurements of echolocation by the harbor porpoise, Phocoena phocoena. He maintains one of the few members of this species in captivity. He has no plans to carry out any behavioral measurements of echolocation in the immediate future, but has done and will continue doing recording and analysis of the sounds emitted by this animal. Professor Andersen has two graduate students from Sweden who intend to examine the signals used by this species for communication. The latter work is contingent upon obtaining a second member of the species.



## **APPENDIX II**

### **TABULATION OF BIOLOGICAL SONAR DATA FOR BATS**

**Preceding page blank**

**TABLE A**  
**TABULATIONS OF BIOLOGICAL SONAR DATA FOR OLD WORLD BATS**

**Preceding page blank**

1  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY			FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^a$ , cm	DURATION, msec	SPL, dB re 0.0005 dyn/cm <sup>2</sup>	FREQUENCY, kHz	$\Delta\lambda^a$ , cm	DURATION, msec	
<i>Aeollia tridens</i>	CF/PN	115-120	~3	6.8		120-95	3-3.6	~1	Major component - 2nd harmonic; pulses produced in groups--groups of 1-10 within 170 msec; instantaneous repetition rates up to 75 pulses/sec
		120	2.9	~9		120-60	2.9-5.8		
		117	2.9			117-95	2.9-3.6		
<i>Aeollicus tricuspidatus</i>	Mixed	112	3.1	1.6-2.4		112-94	3.1-3.7	1.1	Tropical
<i>Eballonura nigrescens</i>	Mixed PN	61 (2nd harmonic)	5.6	0-0.25		61-37 (2nd harmonic)	5.6-9.3	0.6-1.3	Multiple harmonics Tropical
<i>Eptesicus tenuipinnis</i>	PN					62-37	5.5-9.3	0.9-1.9	Temperate
<i>Hipposideros beatus maximus</i>	CF	100-114	3.4-3	~8	~95				Hand held Tropical
<i>Hipposideros brachyotus</i>	CF	85-95	4-3.7	5-8					Hand held Tropical
<i>Hipposideros carteri</i>	CF/PN	140-150 137	2.5-2.3 2.5	6-23		137-100	2.5-3.4		Tropical

<sup>a)</sup> C = 344mm/sec

Preceding page blank

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda$ ( $\mu$ ), m.	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>	FREQUENCY SWEEP, kHz	$\Delta\lambda$ ( $\mu$ ), m	DURATION, msec		
Hipposideros calcaratus	mixed	62 128 (2nd harmonic)	5.5 2.7	3.4-4.6		62-50 128-97 (2nd harmonic)	5.5-6.9 2.7-3.5	~1	No clear evidence that Hipposideros more skillful at echolocation than any other Hipposideros Hipposideros - general correlation between size of bat and frequency used Frequency pattern - principal frequency one octave higher than a very low amplitude component Primarily 2nd harmonic Tropical	Grinnell 1972 Roberts 1972
Hipposideros comersoni	CF/FM	56-68 56 and 66	6.1-5 6.1 and 5.2	9-20		66-57	5.2-6.9	3-3.5	Possibly subspecies differentiation; frequency correlated with body size	Pye and Roberts 1970 Pye 1972 Roberts 1972
Hipposideros cupido	mixed	123	2.8	2.2-2.8		123-97	2.8-3.5	0.8		Grinnell 1972 Roberts 1972
Hipposideros cyclops	CF	101-109	34-32	17-24	~115				Hand held Tropical	Kovick 1956
Hipposideros diadema	mixed CF	58 (2nd harmonic)	5.9	4.8-9.1		58-47 (2nd harmonic)	5.9-7.7	~1	Four harmonics detectable in Hipposideros diadema Typical	Grinnell 1972 Roberts 1972
Hipposideros galeritus		145	2.4	1.9-3.6		145-120	2.4-2.9	~1	Tropical	Grinnell 1972 Roberts 1972

( $\mu$ ) C = 344mm/sec

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY					FM			REMARKS	REFERENCES
		FREQUENCY, MHz	$\lambda^{(a)}$ , mm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>	FREQUENCY SWEEP, MHz	$\Delta\lambda^{(a)}$ , mm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>		
<i>Hipposideros lanthanda</i>	CF	57-74	6-4.6	8-13	~130	Down, 2-8 (55-67)	0.3-0.5			Hand held Tropical	Novick 1958
<i>Hipposideros obecurus</i>	CF	95-109	3.6-3.3	9	~125					Hand held Tropical	Novick 1958
<i>Hipposideros s. speoris</i>	CF	80-120	4.3-2.9	6-10	120-135					Hand held Tropical	Novick 1958
<i>Miniopterus australis paulus</i>	FM					50-17	6.9-20	1.1-2.8		Hand held; 2nd and 3rd harmonics	Novick 1958
<i>Miniopterus fuliginosus</i>	FM					~55-35	6.3-9.8	1.9-3.3	115-120	Hand held	Novick 1958
<i>Miniopterus natalensis</i>	FM					82-50	4.2-6.9	2.7-3.6		Hand held	Novick 1958
<i>Miniopterus schreibersii eschscholtzii</i>	FM					87-40	4.8.6	1.5-3.7	~138	Tropical	Novick 1958
<i>Miniopterus tristis</i>	FM					46-30	7.5-11.5	2.9-6.5	~135	Hand held	Novick 1958

<sup>(a)</sup> C = 344mm/msec

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM			REMARKS	REFERENCES
		FREQUENCY, MHz	$\lambda^{(a)}$ , cm	DURATION, msec	SPL, dB @ 0.002 $\frac{\text{cm}^2}{\text{m}^2}$	FREQUENCY SWEEP, MHz	$\Delta\lambda^{(a)}$ , cm	DURATION, msec		
Lyraderma l. lyra	CF	60-80	5.7-4.3		100-105				Tropical	Novick 1958
Megaderma lyra	CF	58.5 (3rd harmonic) 78.5 (4th harmonic)	5.9 4.4	1.2 (0.4-1.8)					Multiple harmonics Pulses extremely short; varies 1300 msec related to orientational situation. Longer signals are produced in free and unimpeded flight in open space. Shorter (less than 1000 msec) and higher frequency sounds are produced by nonflying bats in exploring environment, or flying bats in approaching a goal or obstacle. Highest frequency - most intense Frequency modulation is nearly absent Tropical	Novick 1962 Schnitzler 1967 Mohr 1967
Megaderma s. spasma	CF	20	17.2						2nd and 3rd harmonics present from beginning at significant amplitudes while 4th harmonic sometimes appeared late in pulse Tropical	Novick 1958

(a)  $C = 344 \text{ m/sec}$

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM				REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^{(a)}$ , cm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>	FREQUENCY SWEEP, kHz	$\Delta\lambda^{(a)}$ , cm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>		
<i>Myotis sacrocarus</i>	FM					50-27	6.9-12.7	2.5-5.7	130	Temperate	Movick 1958
	FM					75-30 (search) 45-25 (terminal)	4.6-11.5	1.5-1.0		Temperate	Slakin 1969
	FM					105-30 (search) 60-30 (terminal)	3.3-11.5	3.5-1.5		Temperate	Slakin 1970
<i>Nycterus arge</i>	CF	20-22 (1st harmonic)	17-15	0.4-1.4	110-120					3-5 Harmonics Band held Tropical	Konstantinov and Alrapetiantz 1968 Slakin 1969 Alrapetiantz and Konstantinov 1967 Alrapetiantz, Konstantinov, and Matjushkin 1969 Konstantinov 1969
	CF	20-22 (1st harmonic)	17-15	0.4-1.4	110-120					3-5 Harmonics Band held Tropical	Movick 1958
	CF	20-22 (1st harmonic)	17-15	0.4-1.4	110-120					3-5 Harmonics Band held Tropical	Movick 1958

(a)  $C = 344 \text{ m/sec}$

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM				REMARKS	REFERENCES
		FREQUENCY, MHz	$\lambda^{(a)}$ , mm	DURATION, msec	SFL, dB re 0.002 $\frac{\text{W}}{\text{cm}^2}$	FREQUENCY SWEEP, MHz	$\Delta\lambda^{(a)}$ , mm	DURATION, msec	SFL, dB re 0.002 $\frac{\text{W}}{\text{cm}^2}$		
Pipistrellus c. ceylonicus	FM					35-25	9.8-13.8	1.7-3.3	95-105*	*At 15 cm from head Tropical	Novick 1958
Pipistrellus coromandra	FM					71-28	4.8-12.3	0.8-3.6	110-115	Band held	Novick 1958
Pipistrellus m. mimus	FM					60-37	5.7-9.3	2.2-4.3		Band held	Novick 1958
Pipistrellus papuanus	FM					75-35	4.6-10	1-2			Grinnell 1972
Rhinolophus alcyon	CF/FM	85-90 89	4-3.8 3.9	11-60		89-69	3.9-5				Pye and Roberts 1970 Roberts 1972
Rhinolophus a. arcuatus	CF/FM	55-65	6.3-5.3	17-34		~66-51	3.2-6.8			Band held	Novick 1958
Rhinolophus euryale	CF/FM	103-104	3.3	35-45 (search) 7-10 (burst)		103-90	3.3-3.8	1.5-3		Flying animals lower the frequency of the CF part by such an amount that the Doppler shifts caused by flight velocity are compensated. Use burst pulse transmission mode during terminal phase of pursuit or landing	Schnitzler 1967 Pye and Roberts 1970

<sup>(a)</sup> C = 344mm/msec



# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY			FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^{(a)}$ , mm	DURATION, msec	FREQUENCY SWEEP, kHz	$\Delta\lambda^{(a)}$ , mm	DURATION, msec		
Rhinolophus ferrus equinus	CP/FM	83.5 (at rest)	4.1	up to 100, 25-50 average (search) 10, burst mode burst duration 30-60	83-56	4.1-6.1	~1.5	Very intense 2nd harmonic of weak fundamental frequency of 41.5 kHz Flying bats lower frequency of CP part to compensate for Doppler shifts due to target or bat motion. Frequency varies between 81.0 and 83.5 kHz dependent on flight velocity, whereas echo frequency is kept constant at 83.5 kHz Uses burst pulse mode of 2 to 10 short pulses/burst during insect capture or obstacle avoidance With decreasing sound duration, duration of FM part diminishes. CP portion continually sacrificed for FM portion, but never completely disappears	Airapetian, Konstantinov, and Metjubkin 1969 Konstantinov and Sokolov 1969 Morick 1971 Mauweller, Schuller, Schnitzler 1971 Schnitzler 1967 Mauweller 1970 Schnitzler 1970 Webster 1963 Simmons 1969, 1971, 1972 Nagus 1967 Pye 1960, 1961 Griffin, Dunning, Cahlander, Webster 1962 Griffin 1962 Morick 1976 A. Pye 1967

<sup>(a)</sup>  $c = 344 \text{ mm/sec}$

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^a$ , mm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>	FREQUENCY SWEEP, kHz	$\Delta\lambda^a$ , mm	DURATION, msec		
Rhinolophus funigatus	CF/FM	45-50 45	7.7-6.9 7.7		15-65	45-35	7.7-10.5			Pye and Roberts 1970 Roberts 1972
Rhinolophus hipposideros	CF/FM	105-115 109	3.3-3 5.2	15-65		109-80	5.2-4.3			Pye and Roberts 1970 Roberts 1972
Rhinolophus landeri	CF/FM	115-122 121	3-2.8 2.8	14-72		121-80	2.8-3.4			Pye and Roberts 1970
Rhinolophus luctus	CF	42	8.2	42-30	8.2-11.5					Roberts 1972
Rhinolophus mehelyi	CF/FM	105	3.3	~35	3.3-3.8	105-90	1.5-3		During pursuit/avoidance uses burst pulse transmission; up to 8 pulses/burst; bursts correlated with respiration	Konstantinov and Sobolov 1969
Rhinolophus r. rouxi	CF	60-75	5.7-4.6	21-45	~130	~65-60	5.3-5.7		Band held	Kovick 1958
Rhinolophus s. subrufus	CF/FM	45 (2nd harmonic) 68-82 (dominant)		26-42 with extremes of 15-60 msec	134	82-51	4.2-6.7			Kovick 1958
Rousettus sp. (Megachiroptera)									Generates clicks by movements of the tongue and exits these through corner of mouth Principal frequencies, 12 and 18 kHz, depending on species and individual; overtones and harmonics present to a considerable degree Tropical	Griffin, Kovick, Hornfield 1958

<sup>a</sup>  $\lambda = 344 \text{ mm/msec}$

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL PATTERN	CONTINUOUS FREQUENCY			FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda$ , m	DURATION, msec	SPL, dB re 0.002 $\frac{\text{W}}{\text{cm}^2}$	FREQUENCY SWEEP, kHz	$\Delta f$ , mm	DURATION, msec	SPL, dB re 0.002 $\frac{\text{W}}{\text{cm}^2}$
<i>Tadarida lucronus</i>	FM					43-25	8-13.8	6.3-10	
<i>Tadarida aildas</i>	FM					22-11	15.5-31	7-10	
<i>Tadarida fragata</i>	FM					28-22.5	12.3-27.5	5.1-8.1	110-115
<i>Tapasous melanopygon</i>	FM					15-12	22.9-28.7	5.1	
<i>Tapasous philip nensis</i>	FM					16-13	21.5-26.5	4.6	
<i>Trisopsa alar</i>	CF/FM	78-92 79 and 88	4.4-3.7 4.4 and 3.9	6-16		~85-65	4-5.3	~3	
<i>Tylonycteris pachypus mayeri</i>	FM					80-55	4.3-6.5	1.5-5	130-145
<i>Vesperugo kuhlii</i>	FM					80-55	4.3-9.8	1-2.2	

(a) C = 344mm/sec

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY			PM			REMARKS	REFERENCES
		FREQUENCY, MHz	$\lambda(m)$ , mm	DURATION, msec	FREQUENCY SWEEP, MHz	$\Delta\lambda(m)$ , mm	DURATION, msec		
Vespertilio murinus	PM				50-80	6.9-17.2	0.6-1.6	Temperate	Slarkin 1970
Vespertilio naebuefi	PM				90-90	3.8-11.4	0.8-2.9	Temperate	Slarkin 1970
Vespertilio nilssonii	PM				50-82	6.9-15.6	2-3	Temperate	Slarkin 1970
Vespertilio ogilvi	PM				75-85	4.6-13.8	1.5	Temperate	Slarkin 1970
Vespertilio savii	PM				80-90	4.3-11.5	0.7-2.0	Temperate	Slarkin 1970
Vespertilio serotinus	PM				60-90	5.7-11.4	1.4-2.2	Has 2nd and sometimes 3rd harmonic component Temperate	Slarkin 1970
Vespertilio superans	PM				50-19	6.9-18.1	1.2-2.5	Temperate	Slarkin 1970

(\*) C = 344mm/msec

## II SONAR CHARACTERISTICS

SPECIES	FLYING SPEED, m/sec	PRF, sec <sup>-1</sup>	CONTINUOUS FREQUENCY		PN		REMARKS	REFERENCES
			BEAMWIDTH, deg	Δ BEAMWIDTH, deg	RANGE RESOLUTION, m Δ r	Δ f		
Megaderma lyra	(see remarks)	10-50 (average) to 500 (peak)	70 (-5 dB) 60 (-5 dB)				When approaching an obstacle, emits pulses in doublets Can hover in flight Assumed dipole source (double emitter, nostrils separated by $\lambda/2$ )	Mogus 1970 Moures 1967 Kovick 1958
Myotis myotis		15-30 40-60 (burst) to 200 (terminal)			~26 ~4 (terminal); 25-70 (search) 8-17 (terminal)	~0.3 ~5	Pulses emitted in packets, or volleys, of 2 to 6 pulses/packet during approach to target	Siakin 1969 Aireprians, Konstantinov, and Stjughin 1969
Myotis myotis		7-10 (search) 40-60 (burst) 100 (terminal)			25-70 (search) 8-17 (terminal)	~0.25 ~0.8	Pulses emitted in packets, or volleys, of 2 to 6 pulses/packet during approach to target	Siakin 1970, 1969 Aireprians, Konstantinov, and Stjughin 1969
Myotis myotis	4-5	5-20 (search) 70-80 (burst) 4-8 (burst rate)	60 (-5 dB) (calculated) (horizontal) 52 (-5 dB) (horizontal) 20 (-1.5 dB) (vertical)		~260 ~10		When investigating unknown object, it emits bursts of shorter pulses, up to 20/burst, at increased pulse repetition frequency, e.g., average PRF during volley of 8 pulses is 35/sec. Pulse duration during burst is ~10 msec. Assumed dipole source (double emitter; nostrils separated by $\lambda/2$ ) Apparent dipole source, both planes	Airapetians, Konstantinov, and Matjushkin 1969 Konstantinov and Sokolov 1969 Schmitzler 1970, 1967 Mogus 1970, 1967 Pye 1960 Griffin, Deamin, Cahlander, Webster 1962 Kovick 1961 Sokolov and Maharov 1971

# **AUDOMETRIC DATA**

SPECIES	AUSCULTATION	FREQUENCY RANGE, kHz	MAXIMUM SENSITIVITY		REMARKS	REFERENCES
			FREQUENCY, kHz	SPL, dB re 0.002 dyn/cm <sup>2</sup>		
<i>Aselliscus tricuspidatus</i>	Electrophysiological (H <sub>1</sub> )	10-150	50-110	25	Roll-offs 10-100 dB/0.01 Δf; single units (neurons), 500-1500 dB/octave	Grinnell 1972
<i>Euballanura nigriscens</i>	Electrophysiological (H <sub>1</sub> )	10-120	50-65	25		Grinnell 1972
<i>Hippodideros calcaratus</i>	Electrophysiological (H <sub>1</sub> )	10-140	50-60 100-130	30		Grinnell 1972
<i>Hippodideros diadema</i>	Electrophysiological (H <sub>1</sub> )	10-90	50-60	30		Grinnell 1972
<i>Hippodideros galericus</i>	Electrophysiological (H <sub>1</sub> )	10-150	60-70 120-140	30-35		Grinnell 1972
<i>Myotis oxygnathus</i>	Behavioral; operant conditioning	0.5-250	40 (10-90)	-15 (40 kHz)	Bats continued to react to all frequencies presented up to limits of generator used--frequency of 250 kHz displaying level of positive responses higher than 70%	Konstantinov, Sanotskaya, and Sobolova 1971 Airepetyans and Konstantinov 1971
<i>Pipistrellus papuensis</i>	Electrophysiological (H <sub>1</sub> )	10-110	40-90	40		Grinnell 1972

UNIVERSAL TALK  
ELECTROPHYSIOLOGICAL

# III AUDIOMETRIC DATA

SPECIES	AUDIOGRAM(1)	FREQUENCY RANGE, kHz	MAXIMUM SENSITIVITY		REMARKS	REFERENCES
			FREQUENCY, kHz	SE, dB re 0.002 dyne/cm <sup>2</sup>		
Rhinolophus euryale	Electrophysiological (N <sub>1</sub> )	<50-150	50-95 104	25-40 ~10	Threshold increases by an average of 6.5 dB/0.1% Δf when signal frequency is lowered by 100 Hz Average threshold slope of 40-55 dB/1.8 kHz	Schnittler, Schuller, and Neuweiller 1971
Rhinolophus ferrum equinum	Electrophysiological (N <sub>1</sub> )	10-100	20-40 and 85.5	0-5		Airapetian, Konstantinov, and Matjushkin 1969 Vasil'ev 1967 Neuweiller 1970 Neuweiller, Schuller, Schnittler 1971
Mussetus amplexicaudatus stresemanni	Electrophysiological (N <sub>1</sub> )	10-100	45-50	50		Grinnell and Magivara 1972

149 BMAVORAL TASK  
ELECTROPHYSIOLOGICAL

**IV**  
**DETECTION/DISCRIMINATION DATA**

SPECIES	TARGET/Obstacle: DETECTION/DISCRIMINATION				SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	SIGNAL FORM	TASK	PERFORMANCE	
<i>Hippodamia diadema</i>						electro-physiological ( $R_n$ )	tone burst	recovery, time interval 50% 100%	3 msec 6-8 msec	Grinnell 1972
<i>Hippodamia galleritius</i>						electro-physiological ( $R_n$ )	tone burst	recovery, time interval 100%	2 msec	Grinnell 1972
<i>Megaderma lyra</i>	nylon monofilament, grid barrier (14 cm separation)	0.008 0.006	avoidance		>50% 18%					Mohr 1967
<i>Myotis myotis</i>	discs squares triangles 6-point stars, aluminum	equal area (disc, 5 diam)	discrimination, shape							Simkin 1969
<i>Myotis myotis</i>	discs squares triangles 6-point stars, aluminum	equal area (disc, 5 diam)	discrimination, shape							Simkin 1969
	multistage pyramid	base, 10x10	detection and recognition							Retraining to small pyramid, signal duration diminished 50% Simkin 1970

<sup>(1)</sup> BINAURAL: TALK ELECTROPHYSIOLOGICAL



IV  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION					SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(c)</sup>	SIGNAL FORM	TASK	PERFORMANCE		
Myotis otis (Cont'd)	small multistage pyramid	base 4.4x4.4	detection and recognition, selection in presence of large pyra- mid and sphere							Identification of spheres, uses long, high frequency signals	
	spheres	4.4 and 2.9	detection and recognition, selection in presence of pyramid and other spheres							Approached positive shape alone. Did not approach negative shape alone	Konstantinov and Abbasova 1968
	squares versus triangle and disc	equal area (156 cm <sup>2</sup> )	discrimina- tion, shape	160-360	~9%						
	squares (6)	area, 156 cm <sup>2</sup> to 625 cm <sup>2</sup>	discrimina- tion, size	160-360	99%, 156 cm <sup>2</sup> versus 625 cm <sup>2</sup> 80%, 156 cm <sup>2</sup> versus 172 cm <sup>2</sup> 87%						
	squares, aluminum, versus square, plywood	area 156 cm <sup>2</sup>	discrimina- tion, material	160-360							
	metal wire barrier	0.008	avoidance			behavioral		discrimina- tion, target range difference 2-1 m	77%, 20-2.5 cm (20:160 msec)	Calculated echo level, -1.5 dB re 0.0002 dynes/cm <sup>2</sup>	Airapetians and Konstantinov 1971
	cylinders	20 long									

(c) BEHAVIORAL; TASK  
ELECTROMYOLOGICAL

II  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION				SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(c)</sup>	SIGNAL FORM	TASK	PERFORMANCE	
Myotis oxyanthus (Cont'd)	squares, aluminum brass plexiglass	1x1x1	discrimina- tion: edge smooth versus serrated contour plane versus convex material metal versus metal metal versus other		70%-100%  70%-100%  50% 70%-100%					Airapetians, Konstantinov, and Matjuhakin 1969
Myotis noctula	cube cylinder pyramid, plastic	equal volumes	discrimina- tion, form	30-50						Airapetians and Konstantinov 1965
Pipistrellus pauanus						electrophys- iological (M <sub>1</sub> )	tone burst	recovery, time inter- val 50% 100%	1-1.5 msec 2 msec	Grinnell 1972
Plecotus auritus	cube cylinder pyramid, plastic	equal volumes	discrimina- tion, form	50-50						Airapetians and Konstantinov 1965
Rhinolophus euryale	metal wire barrier (17.5 cm separation)	0.02 0.008	avoidance		70% 68%					Schnittler 1967

(c) BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL

II  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/DISTANCE: DETECTION/DISCRIMINATION				SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE (C)	SIGNAL FORM	TASK	PERFORMANCE	
Rhinolophus ferrum-equinum	cylinders	20 long				behavioral		discriminate target range difference; R=1 m	75%; ΔR=4 cm (Δt=250 μsec)	Alrapetianz and Konstantinov 1971
	cylinders	20 long				behavioral		discriminate crossrange target separation; R=2.5 m	4°30'	
	squares, aluminum brass plexiglass	1x1x1	discrimination: edge smooth versus serrated contour plane versus convex material metal versus metal metal versus other		70%-100%					Alrapetianz, Konstantinov, and Matyushkin 1969
					70%-100%					
					50%					Simmons 1972
					70%-100%					
					70%-100%					Schnitzler 1967
					70%-100%					
	metal wire barrier (17.5 cm separation)	0.02 0.008	avoidance		70% 50%	behavioral, artificial echo	tone burst	compensate for simulated target velocity (Doppler)	ΔR 0.2 m/sec (~20 km/h) 80 km/h	Möhres 1960
	square	20x20	recognition	1000						

(C) BEHAVIORAL: TASK  
ELECTROMYOLOGICAL

**II**  
**DETECTION/DISCRIMINATION DATA**

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION				SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(c)</sup>	SIGNAL FORM	PERFORMANCE		
Rhinolophus schelyi	metal wire barrier (50 cm separation)	0.008	avoidance		~75%-80%					Konstantinov, Sokolov, and Stosman 1967
Rousettus aegyptiacus (-megachiroptera)	metal wire barrier (53 cm separation)	0.15 diam 0.107 diam	avoidance		77% 68%				Are more vulnerable to noise interference than Vespertilionidae	Griffin, Novick, and Kernfield 1958

(c) BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL

Y  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Asellia tridens</i>		desert, palm groves, Old World	Fly swiftly near ground			Walker 1968
<i>Hipposideros</i> sp.		paleotropical	Flv close to ground. Hk insects in flight while following contours of ground and vegetation	insects (beetle, termites, cicadas) fruit		Movick 1958 Walker 1968
<i>Megaderma</i> sp.	20-50 g	Feed among trees and undergrowth. paleotropical	Adept flyers. Can hover in flight. Hunt along solid surfaces	sitting or crawling small insects and vertebrates; carnivorous	Heavily reliant upon vision under normal circumstances	Mohres 1967 Movick 1958 Walker 1968
<i>Miniopterus</i> sp.	8-20 g	tropical and temperate zones of Old World	Rapid, jerky flight. Feed at altitudes of 10-20 m	insects		Movick 1958 Walker 1968
<i>Myotis macrotarsus</i>				fish and insects		Movick 1958
<i>Myotis oxygnathus</i>	6-7.5 cm long		High maneuverability in flight	insects		Konstantinov and Abbasova 1968
<i>Myotis</i> sp.	to 30 g	rain forest to arid plains paleotropical	Hunt insects on solid surfaces	insects (moths, beetles)		Movick 1958 Walker 1968

Y  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Nyctalus noctula</i>	15-40 g	tropical and temperate zones of Old World	High maneuverability in flight	small animals-- rodents winged insects-- beetles, ants, moths		Gould 1975 Walher 1968
<i>Pipistrellus</i> sp.	~4-8 g	worldwide	Jerky flight	insects (moths, mosquitoes, gnats)		Walher 1968
<i>Placotus auritus</i>	5-20 g	temperate regions of Old World	Hover in flight. Hunt insects on solid surfaces	insects (moths, beetles)		Novick 1978 Gould 1975 Walher 1968
<i>Rhinolophus</i> sp.	to 28 g	Tropical and temperate zones of Old World. Prefer open country for hunting. Feed near and on ground	Flight may be swift, straight, and sometimes high	insects (moths, beetles, spiders)		Novick 1978 Schnitzler 1970 Walher 1968
<i>Tapbazous</i> sp.	10-30 g	open areas paleotropical		insects		Novick 1978 Walher 1968
<i>Tylonycteris pachypus</i>	smallest known bat	paleotropical		insects		Novick 1978

Y  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
Vespertilio kuhlii*		open country	Fly near ground	insects and fish (mosquitoes)	*Walker lists only two species of the genus Vespertilio: V. murinus and V. superans, found in temperate regions of Old World	Sinkin 1970 Walker 1968
Vespertilio murinus	14 g	forests and river valleys	Fly high above ground	insects and fish (mosquitoes, beetles, moths, butterflies)		
Vespertilio nathusii		temperate forest areas and water courses	Fly high above ground	insects and fish (mosquitoes)		
Vespertilio nilsonii		near forest areas		insects and fish (mosquitoes)		
Vespertilio ognevi				insects and fish		
Vespertilio savii		grassy plains, along water courses		insects and fish (mosquitoes)		
Vespertilio serotinus		open country, sand dunes	Fly near ground. Can hover in flight	insects and fish (beetles)		
Vespertilio superans				insects and fish (mosquitoes)		

**TABLE B**  
**TABULATIONS OF BIOLOGICAL SONAR DATA FOR NEW WORLD BATS**

**Preceding page blank**



I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY			PM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^{(a)}$ , mm	DURATION, msec	SPL, dB re 0.0002 $\frac{\text{W}}{\text{cm}^2}$	FREQUENCY, kHz	$\Delta\lambda^{(a)}$ , mm	DURATION, msec	SPL, dB re 0.0002 $\frac{\text{W}}{\text{cm}^2}$
Anousa geoffroyi	FM					~100-50	5.4-6.9	~0.5-2	
Artibeus cinereus	FM					90-60	3.8-5.7		
Artibeus jamaicensis palmarum	FM					56-51	6.1-11.1	2.7-3.2	95
						65-42	5.3-8.2	1-3	
Artibeus lituratus	FM					~5-40	5.1-8.6		
Malantiopteryx plicata	FM					18-15	18.9-22.5		2-4.2
Carollia perspicillata	FM					80-55 (2nd harmonic)	4.3-6.5	0.5-1	
Centurio senex	FM					115-70	3-4.9	2	
Chilonycteris personata mexicana	FM					55-38	6.5-9.1	~3.2	

(a)  $C = 344 \text{ m/sec}$

Preceding page blank

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^{(0)}$ , cm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>	FREQUENCY SWEEP, kHz	$\Delta\lambda^{(0)}$ , cm	DURATION, msec		
Chilonycteris pallotis	Mixed	~21 (1st harmonic)	16.4	3-5 (search) 0.6-1 (terminal)		21-17	16.4-20	~1	1st and 2nd harmonics always present Maintains constant pulse-echo overlap (~1 msec) during pursuit FM occurs midway in pulse Tropical	Movick 1966 Movick 1971
Chilonycteris rubiginosa (chilonycteris parnellii)	CF	32 (1st harmonic) 64 (2nd harmonic)	10.8 5.4	14-26 (search) 9 (terminal)		32-28 64-56	10.8-12.5 5.4-6.7	1.5-2	2nd harmonic dominant At rest--maintains frequency within $\pm 100$ Hz of average frequency In flight--lowers frequency to compensate for Doppler shifts caused by flight velocity. Appears to detect insects initially by pulse-echo overlap Following detection, pulse duration increases until pulse-echo overlap exceeds 17-18 msec. Uses burst pulse transmission mode during pursuit or landing; up to 30 pulses/burst pulse duration: $\frac{1}{2}$ msec Tropical	Movick 1971 Movick 1966 Movick and Vainys 1964 Grinnell 1970 Griffin and Movick 1975 Mogus 1967 A. Pye 1967 Schnitzler 1970
Chiroderma villosus	FM					102-60	3.4-5.7	1.0-1.5	Signal contains several harmonics Tropical	A. Pye 1967
Desmodus rotundus aurinus	FM					75-48 (2nd harmonic)	4.6-7.2	0.8-1.6	2nd harmonic present at high amplitude Tropical	Movick 1963 A. Pye 1967 Griffin and Movick 1975

(a) C = 344m/sec

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY			FM			REMARKS	REFERENCES
		FREQUENCY, MHz	$\lambda^{(1)}$ , m	DURATION, msec	FREQUENCY SWEEP, kHz	$\Delta\lambda^{(1)}$ , m	DURATION, msec		
Eptesicus fuscus	FM				55-20 (in laboratory)		1-15 (1-5 in laboratory) (10-15 in open)	Energy peak - 30 kHz Indoors - pulses lasting 4 msec or less; flying in straight lines at several meters above the ground their pulses almost always lasted 10 msec or longer Out of doors - pulses are CF during major part of duration; frequency change only, 8.2 kHz over 8.6 msec (outdoors) and drops from 50-25 kHz during 2 msec (laboratory) Temperate	Peir and Simmons 1971 Simmons and Vernon 1971 Simmons 1971, 1969 Mogus 1967 Griffin 1962, 1958, 1953 Webster and Brazier 1968
Glossophaga longirostris	FM				112-56	3.1-6.1	0.5-2	Tropical	A. Pye 1967
Glossophaga soricina leachii	FM				95-60	3.7-5.7	0.9 (0.7-1)	Strong 2nd harmonic	Novick 1965 Griffin and Novick 1955
Lasiurus borealis	FM				95-40 (search)	3.4-8.6	3-0.5 (in laboratory)		Webster and Brazier 1968 Webster 1965 Griffin 1955
Leptonycteris nivalis	FM				95-35 (terminal)	6.2-10	6-11 (in open)	Tropical	Novick 1965
Lonchophylla robusta	CF	25-108	13.8-3.2	0.5-2.7	100-50 (2nd harmonic)	3.4-6.9	2-8	Many harmonics Tropical	Griffin and Novick 1955

<sup>(1)</sup> C = 344m/sec

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				PM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^{(a)}$ , mm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>	FREQUENCY SWEEP, kHz	$\Delta\lambda^{(a)}$ , mm	DURATION, msec		
Lonchorhina aurita	CF	12 kHz	28.7	1-5.5					2nd and 3rd harmonics very prominent Tropical	Griffin and Novick 1955
Macropyllus macropyllus	CF	21-30	16.4-11.5	0.9-1.9			8.6-13.2	2.5-5.5		Griffin and Novick 1955
Macrotus mexicanus	FM					40-26			Signal accompanied by strong 2nd harmonic Tropical	Novick 1965
Mormonops megalophylla	FM					40-37	7.2-9.5	4-5	Tropical	A. Pye 1967
Myotis lucifugus	FM					100-40 (search) 35-25 (terminal)	3.4-8.6 9.8-13.6	3-5 0.5-0.1	95-115 Temperate	Griffin 1958 Novick 1971 Griffin 1962 Webster and Brazier 1988
										Cablander 1967 Mogus 1967
Natalus mexicanus	FM					89-50	4-6.9	2.3-2.5	Strong 2nd harmonic Tropical	Novick 1962
Neotoma labialis minor (Diras albiventer minor)	FM					62-41	5.5-8.4	4.1-13	Tropical	Griffin and Novick 1955

<sup>(a)</sup> C = 344mm/msec

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^0$ , mm	DURATION, msec	SPL, dB re 0.0002 dyn/cm <sup>2</sup>	FREQUENCY SWEEP, kHz	$\Delta\lambda^0$ , mm	DURATION, msec		
Nocilio leporinus	Mixed CF	60	5.7	3.7	125	60-30	5.7-11.4	93.7	Captive, flying in outdoor cage +1 msec terminal	Suthers 1965
	Mixed CF	60	5.7	6.9-8.9 7.1 13.8				7.1	Wild - duration twice that of caged Emits pulses in triplets: mixed, CF, mixed Emits CF in terminal phase of capture Does not shorten pulse during wire avoidance Tropical	Mogus 1967 Griffin 1958 Suthers 1967 Griffin and Novick 1955 Pye 1966
Phylloderma stenops	FM					70-35	4.9-9.8	0.8-2	Tropical	A. Pye 1967
Phyllotomus discolor	FM					50-25	6.9-13.8	0.5-4	Multiple harmonics Tropical	A. Pye 1967
Phyllotomus hastatus panamensis	FM					65-25	5.3-13.8	0.5-4	Several overlapping, harmonically related sweeps Energy peak of 35-40 kHz Tropical	Paff and Simmons 1971 Simmons 1971 Grinnell 1970 Griffin and Novick 1975 A. Pye 1967
Piconyx vivax	FM					40-20	8.6-17.2	3	Pulses recorded during flight relatively constant in frequency, from 20-27 kHz Duration does not normally shorten significantly with increasing FFR Temperate	Suthers 1967 Webster and Brazier 1969

(\*) C = 344mm/sec

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY				FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda^0$ , cm	DURATION, msec	$\frac{SPL}{0.002 \frac{m}{sec}}$	FREQUENCY SWEEP, kHz	$\Delta\lambda^0$ , cm	DURATION, msec	$\frac{SPL}{0.002 \frac{m}{sec}}$	
<i>Plecotus rafinesquii</i>	FM					80-85 (search ?)	4.3-13.8	4	85	Bats from nostrils or mouth
						45-85 (terminal?)	7.6-13.8	1		Temperate
<i>Plecotus townsendii</i>	FM					80-20	4.3-17.1	2-5 (search)	60-70	Temperate
								0.3-0.5 (terminal)		
<i>Pteronotus davyi</i>	Mixed	59 (1st harmonic)	8.8	3.9-5 (search)		39-31.5	8.8-10.5	3.4 (search)		Strong 2nd harmonic
		78 (2nd harmonic)	4.4	1-1.25 (terminal)		78-63	4.4-5.5	1.05-1.0 (terminal)		Tropical
<i>Pteronotus swainsonii</i>	CF	~25	13.8	2.2						
		52 (2nd harmonic)	6.6	1.5			6.6-8.2	1.5		2nd-5th harmonics
<i>Rhynchiscus naso</i> ( <i>Rhynchonycteris naso</i> )	CF	90-94 (2nd harmonic)	3.8-3.7	4-6.8						Tropical

$60^\circ C = 344 \text{ m/sec}$

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	CONTINUOUS FREQUENCY			FM			REMARKS	REFERENCES
		FREQUENCY, kHz	$\lambda(\mu)$ , mm	DURATION, msec	SPL, dB re 0.0002 $\frac{\mu W}{cm^2}$ @ 10'	FREQUENCY, kHz	$\Delta\lambda(\mu)$ , mm	DURATION, msec	
Saccopteryx bilineata	CF	20-35	17.2-10.4	3-5-15		20-18 (1st harmonic)	17.2-19.1	8	Multiple harmonics. 2nd dominant Tropical
	FM					40-56 (2nd harmonic)			
Sturnira lilium	FM					96-50	3.6-6.9	1-1.3	Tropical
	FM					96-50	3.6-6.9	1-1.3	
Tadarida brasiliensis mexicana	FM					45-25	7.6-13.8	1.4-4.2	Bats pulse doublets during approach to obstacle
	FM								
Uroderma bilobatum	CF	69-88	5-4	1-2					Tropical
	FM					108-72	3.2-4.8		
Vampyrus belleri	FM					110-65	3.1-5.3	1.8-1.5 (search) 0.5 (terminal)	Tropical
Vampyrus spectrum	FM								Bradbury 1970

(a)  $C = 344 \text{ mm/msec}$

## II SONAR CHARACTERISTICS

SPECIES	FLYING SPEED, m/sec	PRR, sec <sup>-1</sup>	CONTINUOUS FREQUENCY		FM		REMARKS	REFERENCES
			BEAMWIDTH, deg	Δ BEAMWIDTH, deg	RANGE RESOLUTION, m	Δ f		
<i>Chilonycteris parnellii mexicana</i>	4-5	~14 (search) 80-100 (terminal)			~33	~1	Increases pulse duration during early pursuit to increase pulse-echo overlap to ~17 to 20 msec. Decreases pulse duration during later pursuit to reduce pulse-echo overlap to ~5 msec at capture	Novick and Vaisnys 1965 Novick 1965
<i>Chilonycteris peilotis</i>	1.75	17-18 (search) ~170 (terminal)					Maintains 1-1.5 msec pulse-echo overlap during pursuit	Novick 1965, 1963
<i>Chilonycteris rubiginosa</i> ( <i>chilonycteris parnellii</i> )	3	to 125	22 (-3 dB) (calculated) (2nd harmonic)				Assumed simple piston radiator in infinite baffle; 1-56 kHz	Schnitzler 1968 Mogus 1970
<i>Eptesicus fuscus</i>	3-4	4-10 (search) 200 (terminal)	55-65 (-3 dB) (search) 42 (-3 dB) (calculated)		~58 (search) 7.5 (terminal)	0.4-0.8 3-5	Assumed simple piston radiator	Peff and Simmons 1971 Mogus 1967, 1970 Webster and Brasier 1968
<i>Lasurus borealis</i>	4-12	5-15 (search) to 240 (terminal)			40-50 (search) 5-8.5 (terminal)	~0.3 0.5		Webster and Brasier 1969, 1968 Webster 1965
<i>Myotis lucifugus</i>	0.5-3 (search/pursuit) 0.5-1 (capture)	10-20 (search) 100-200 (terminal)	25-50 (-3 dB) (search; calculated) ~70 (-3 dB) (terminal; calculated)		>34 (search) 5-17 (terminal)	~0.4 3-5	Tend to reduce average flight speed in laboratory	Griffin 1958 Webster and Brasier 1968, 1965 Mogus 1970, 1967 Cahlander, McCue and Webster 1964



## II SONAR CHARACTERISTICS

SPECIES	FLYING SPEED, m/sec	PER, sec <sup>-1</sup>	CONTINUOUS FREQUENCY		FM		REMARKS	REFERENCES
			BEAMWIDTH, deg	Δ BEAMWIDTH, dB	RANGE RESOLUTION, m	Δ f		
Noctilio leporinus	5-8	10-50 (search) 200 (terminal)	20 (-3 dB) (calculated)	20-34 (-3 dB) (calculated)	17 (terminal)	0.6	Assumed simple piston radiator in infinite baffle; f=50 kHz; Δf=50-50 kHz	Suthers 1965 Bloedel 1955 Mogus 1967, 1970 Griffin 1958
Phyllostorus hastatus parvulus				~45 (-3 dB)	8-17	~0.5		Peff and Simmons 1971 Simmons 1971
Pisonyx yvesi		10-20 (search) 50-200 (terminal)			~50	0.7	High FMR's unusual; average 50 to 50	Suthers 1967 Webster and Brazier 1969
Macrotus ranfinesqui	(see remarks)	15 (search) ~65 (terminal)			68 17	~0.8	Can hover in flight	Mogus 1967 Webster and Brazier 1963
Pteronotus davyi	1-25	~10 (search) 200 (terminal)			55-85 (search) ~17 (terminal)	2-1	Signal has strong 2nd harmonic Maintains 1-3 msec pulse-echo overlap during pursuit	Horvick 1965, 1963

### III AUDIOMETRIC DATA

SPECIES	AUDIOGRAM(s)	FREQUENCY RANGE, kHz	MAXIMUM SENSITIVITY		REMARKS	REFERENCES
			FREQUENCY, kHz	SPL, dB re 0.0002 dyne/cm <sup>2</sup>		
<i>Carollia perspicillata</i>	Electrophysiological ( $H_k$ )	10-150	80	~25		Grinnell 1970
<i>Chilonycteris parnellii</i>	Electrophysiological (CN)	25-75	61.8		Roll-offs of 150-200 dB/kHz	Pollak, Henson, and Novick 1970
<i>Chilonycteris rubiginosa</i>	Electrophysiological ( $H_k$ )	10-130	63	5	Sharply tuned to frequencies in 2nd harmonics of orientation pulses from 62-65 kHz down to ~55 kHz. Roll-offs of 30 dB/0.2 kHz, 55 dB/0.5 kHz	Grinnell 1970
<i>Desmodus rotundus murinus</i>	Electrophysiological (CN)	n. 1-100	10-60		Removal of pinna and tragus had negligible effect upon response or sensitivity	Vernon and Peterson 1966
<i>Myotis fuscus</i>	Behavioral; Operant conditioning	2.5-100	10-25 50-70	~5	Excellent agreement between behavioral and electrophysiological ( $H_k$ ) measurements of audiogram	Dalland 1965 Dalland, Vernon, and Peterson 1967 Henson 1971
<i>Molossus milleri</i>	Electrophysiological ( $H_k$ )	<10-60	35-40			Henson 1967 Henson 1971

(1) BEHAVIORAL; TASE  
ELECTROPHYSIOLOGICAL

### III AUDIOMETRIC DATA

SPECIES	AUDIOLOGICAL	FREQUENCY RANGE, MHz	MAXIMUM SENSITIVITY		REMARKS	REFERENCES
			FREQUENCY, kHz	SPL, dB 7 0.001 $\mu\text{W}/\text{cm}^2$		
Myotis lucifugus	Electrophysiological (CN)	0.7-100	12 and 35 (12-60)			Vernon, Dalland, and Meyer 1966
	Electrophysiological ( $M_n$ )	10-170	30-50	~25		Grinnell 1965
	Behavioral; operant conditioning	12-120	40 (20-60)	10		Dalland 1965
Phyllostomus hastatus panamensis	Electrophysiological ( $M_n$ )	10-90	30-35 50-55	~25		Grinnell 1970
Plecotus townsendii	Electrophysiological ( $M_n$ )	10-100	15-25 50-65	0-5	Two regions of equally low threshold, one near or slightly below the range of fundamental emitted frequencies, and a second at the second harmonic of the orientation sounds	Grinnell 1965
Pteronotus susupensis	Electrophysiological ( $M_n$ )	10-140	45-55	10-20		Grinnell 1970
Saccopteryx bilineata	Electrophysiological ( $M_n$ )	10-120	40-45	5-20	Roll-offs 10-100 dB/0.01 $\Delta f$ ; single units (neurons), 500-1500 dB/octave	Grinnell 1970
Tadarida brasiliensis mexicana	Electrophysiological (CN)	0.7-100	10-40			Kenson 1971, 1967
Tadarida molassa	Electrophysiological ( $M_n$ )	0.1-45	11-17			Kenson 1971, 1967

(C)BIOLOGICAL: TASK  
ELECTROPHYSIOLOGICAL

**IV**  
**DETECTION/DISCRIMINATION DATA**

SPECIES	TARGET/Obstacle; DETECTION/DISCRIMINATION			SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(c)</sup>	SIGNAL FORM		
<i>Artibeus jamaicensis palmarum</i>	metal wire barrier (50 cm separation)	0.0175 diam	avoidance		72%				Griffin and Novick 1975
<i>Carollia perspicillata</i>	metal wire barrier (20 cm separation)	0.0175 diam	avoidance		56%				Griffin and Novick 1975
<i>Chilonycteris parnellii mexicanus</i>	insects, fruitflies	0.2 (wing-span)	detect/capture	3.8				Assumed detection range based on calculated pulse-echo overlap	Novick and Vainys 1964 Novick 1965
<i>Chilonycteris pusillus</i>	insects, fruitflies	0.2 (wing-span)	detect/capture	40-70				Assumed detection range based on calculated pulse-echo overlap	Novick 1965
<i>Eptesicus fuscus</i>	triangles, plastic	7-10, base 5.5-5, height 68x12" versus 128x68"	discrimination, size	30	75% @ $\Delta$ area = 17% $\Delta$ intensity ~ 1.5-3 dB > 80%			Discrimination likely mediated by echo intensity difference (proportional to $\Delta$ area)	Simmons and Vernon 1971
	triangles, plastic	10 cm, base 5 cm, height 128x68"	discrimination, shape	30					
	triangles, plastic	10 cm, base 5 cm, height 128x68"	discrimination, distance	30, 60	75%, $\Delta t = 1.2$ cm ( $\Delta t: 70-75$ $\mu$ sec)			Resolution apparently independent of distance. Performance compares favorably with estimate based on autocorrelation of emitted signal	Simmons and Vernon 1971 Simmons 1971
	pebbles	1-2 diam	detection	200					Griffin 1975
	spheres, nylon	1.9 diam	horizontal angle resolution	30	75%, $\Delta \theta = 5-8^\circ$				Poff and Simmons 1971

<sup>(c)</sup> BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL

IV  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE: DETECTION/DISCRIMINATION				SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE(s)	SIGNAL FORM	TASK	PERFORMANCE	
Glossophaga	metal wire barrier (50 cm separation)	0.0175	avoidance		80%					Griffin and Kovick 1955
Lasurus borealis									Seen able to evaluate position of maneuvering target without apparent head following	Webster 1965
Neotoma mexicanus	metal wire barrier (40 cm separation)	0.038 diam 0.027 diam	avoidance		80% 77%					Grummon and Kovick 1965
Myotis lucifugus	sphere, nylon mealworm	0.16 0.32 0.2-0.3 diam 1.5-2.5 long	detection discrimina- tion (versus spheres)	~ 60 90-110	30% versus spheres of com- parable echo intensity; ~55% versus others				Are able to select individual target out of cluster (nS16)	Webster and Brasier 1965
	metal wire barrier (50 cm separation)	0.3 0.107 0.065 0.046 0.028 0.018	detection avoidance	215 185 150 120 105 90	80% 75% catches: 90% mealworms versus 20-40% discs				Are capable of achieving localization accuracies of 1 cm <sup>3</sup>	Grinnell and Griffin 1958
	discs, bachelite	1.6 1.25 diam 0.3 thick	discrimina- tion (versus mealworms)							Griffin, Friend, and Webster 1965

(\*) BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL

# IV DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION				SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(a)</sup>	SIGNAL FORM	TASK	PERFORMANCE	
Myotis lucifugus (Cont'd)	disks, metal	0.63 0.95 1.25 1.6 dian 0.05 thick	discrimination (versus mealworms)		catches: 80% mealworm versus 30%, 0.95 20%, 1.25					Griffin, Friend, and Webster 1965
	insects, fruitflies	0.5-3 mg	detection, capture	50-100						Griffin, Webster, and Michael 1960
	mosquitoes	~2.2 mg								
	metal wire barrier (45 cm separation)	0.054 0.028	avoidance		~88% ~88% (in quiet and in noise)					Griffin 1958
	metal wire barrier	0.121 0.026 0.012	avoidance		82% 52% 35%	electrophysiological	tone burst	discrimination, frequency	Δf: 0.5-1 kHz, f: 50-100 kHz	Curtis, 1952
						electrophysiological	tone burst	discrimination, intensity	Δf: 0.2-0.5 dB	Grinnell 1965
						electrophysiological	tone burst	discrimination, time	Are capable of complete temporal resolution down to intervals <1 msec	

<sup>(a)</sup> BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL

IV  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION			SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	SIGNAL FORM	TASK	PERFORMANCE
Noctilio leporinus	cube, fish mus- cle	1-2	detection of cube par- tially exposed at water sur- face	130-150	~100%				
	upwelling, water	5-10, high 10-15 diam	detection and dipping for fish cube		85%				
	metal wire	0.055 0.021	detection of 0.5 cm length pro- truding above water surface	~60	90% 80%				
	metal wire	0.13 and 0.09	discrimina- tion of single 0.13 cm wire from pair of 0.09 cm wires pro- truding 0.05 cm above water surface		~80%				
	metal wire barrier (55 cm separation)	0.051 0.021	avoidance	150 130	76% 60%				
								Are extremely skillful at detecting echoes from tiny surface disturbances	Suthern 1965
								Emitted longer pulses when approaching 0.021 cm wires than when approaching 0.051 cm wires	Suthers 1967

<sup>(1)</sup> BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL

# IV DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE, DETECTION/DISCRIMINATION					SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION			REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	SIGNAL FORM	TASK		
Phyllotis bassatus peruanensis	triangles, plastic	10, base 5, height	discrimina- tion, target distance	60	Ad, ~1.2 cm (75%)				Echo level: 0.05 dyne/cm <sup>2</sup> Performance compares favorably with estimate based on auto- correlation of emitted signal	Simmons 1971
	spheres, nylon	1.9 diam	horizontal angle resolution	30	4°-6°					Peff and Simmons 1971
Pizonyx virgatus	metal wire barrier (55 cm separa- tion)	0.051 0.021	avoidance		85% 51%				Average distance of detection: 0.051: 110 cm 0.021: 70 cm	Suthers 1967
	metal wire barrier (45 cm separa- tion)	0.054 0.028	avoidance		91% 69%					Griffin 1958
Plecotus townsendii (Plecotus gratiosus)	metal wire barrier (45 cm separa- tion)	0.028	avoidance		80% (in quiet) 50-60% (in noise)				Estimated echo level @ 10 cm from wire: 24 dB re 0.0002 dyne/cm <sup>2</sup> Noise SPL: +80-90 (15-55 kHz)	Griffin and Grinnell 1958
	metal wire barrier (45 cm separa- tion)	0.054	avoidance		~85% (in noise; 10-50 kHz bandwidth) 30-40% (in noise; 10-90 kHz bandwidth)				Indicates use of higher (2nd) harmonic of signal (to over- come noise masking)	Griffin, McGue, and Grinnell 1965
						electrophys- iological (N <sub>1</sub> )	tone burst	detection, intensity difference	ΔI: 0.2-0.5 dB	Grinnell 1965
						electrophys- iological (N <sub>1</sub> )	tone burst	detection, frequency difference	ΔF: 0.005f to 0.01f	
						electrophys- iological (N <sub>1</sub> )	tone burst	detection, time difference	Δt: <1 msec	

(1) BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL



**IV**  
**DETECTION/DISCRIMINATION DATA**

SPECIES	TARGET/OBSTACLE: DETECTION/DISCRIMINATION				SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	SIGNAL FORM	PERFORMANCE		
Pteronotus davyi	insects, fruitflies	0.2 (wing- span)	detect/ capture	10-85					Assumed detection range based on calculated pulse-echo over- lap	Novick 1965 Novick 1965
Vampyrus spectrum	spheres, rubber	15 and 6, diam	discrimina- tion, shape		~85%, "x" cm sphere versus cylinder					
	cylinder, wood	10.5 L x 2.5 diam			~80%, 6 cm sphere versus cylinder					Bradbury 1970
	sphere, lucite spheroid, lucite	5.08 diam ratio 5/3, inter- focal dis- tance: 2.1	discrimina- tion, shape	50-150	60%-80%				One bat used frequency responses of targets for discrimination; other used overall amplitude differences	

<sup>(1)</sup>RELAY/MORAL: TASK  
ELECTROPHYSIOLOGICAL

Y  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Anoura geoffroyi</i>	14 g	forest (tropical and arid) neotropical		fruit and insects		Pye 1967 Walber 1968
<i>Artibeus jamaicensis palmarum</i>	55-70 g	neotropical		fruit		Novick 1965 Griffin and Novick 1975 Pye 1967 Griffin 1978 Walber 1968
<i>Artibeus lituratus</i>	45-90 g	neotropical		fruit, nuts		Walber 1968
<i>Elanopteryx plicata</i>	5-9 g	arid to semitropical zones of New World	Slow, erratic flight	insects		Walber 1968
<i>Carollia perspicillata</i>	14-20 g pinnae: 17 mm long 8 mm wide	neotropical	Swift fliers	fruit		Griffin and Novick 1975 Grinnell 1962 Walber 1968
<i>Centurio senex</i>	17-28 g	neotropical	Jerky flight	fruit		Pye 1967 Walber 1968

V  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Chilonycteris personata</i>	~20 g	neotropical		insects		Novick 1963 Griffin 1958 Walker 1968
<i>Chilonycteris rufiginosa</i> ( <i>Chilonycteris parnellii</i> )	18-24 g wings: 22 mm long 9 mm wide	rain forest neotropical		insects		Novick 1963 Griffin and Novick 1955 Pye 1967 Grinnell 1962 Walker 1968
<i>Chiroderma villosus</i>	to 30 g			fruit		Pye 1967
<i>Desmodus rotundus murinus</i>	15-50 g pinnae: 15.5 mm long 9 mm wide	neotropical	Slow, silent flight near ground	Large, not very elusive prey. Feeds exclusively on vertebrate blood		Novick 1963 Vernon and Peterson 1966 Pye 1967 Grinnell 1962 Walker 1968

Y  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Eptesicus fuscus</i>	12-24 g pinnae: 14.5 mm long 8 mm wide		Rapid, strong, steady flight. Feed near ground	Insects		Grinnell 1962 Walker 1968
<i>Glossophaga</i> sp.	8-15 g	wooded country and arid lowlands	Fly at great speed. Hover to drink nectar	fruit and insects		Pye 1967 Griffin and Novick 1975 Walker 1968
<i>Lasiurus borealis</i>	6-30 g	over woods and along water courses	Strong fliers. Feed 6-15 m above ground	Insects (moths)		Gould 1955 Walker 1968
<i>Leptonycteris nivalis</i>	18-30 g	arid, open country		fruit		Novick 1965 Walker 1968
<i>Lonchophylla robusta</i>	45-60 g	neotropical		fruit, nectar, insects		Walker 1968
<i>Lonchorhina aurita</i>	10-16 g	neotropical				Griffin and Novick 1975 Walker 1968
<i>Macrophyllum macrophyllum</i>	6-9 g	rain forest neotropical				Griffin and Novick 1975 Griffin 1976 Walker 1968

Y  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Macrotus mexicanus</i>	12-20 g pinnas: 26 mm long 15 mm wide	arid lowlands	Feed in flight on ground	insects		Novick 1963 Grummon and Novick 1963 Grinnell 1962 Walker 1968
<i>Microonycteris hirsuta</i>	12 g	neotropical		fruit		Pye 1967 Walker 1968
<i>Mollosus milleri</i>	to 95 g	neotropical		insects		Walker 1968
<i>Mormoops megalophylla</i>	12-20 g	hunt over land and water just above surface		insects		Pye 1967 Walker 1968
<i>Myotis lucifugus</i>	5-14 g pinnas: 14 mm long 7 mm wide	wooded and open areas worldwide	Erratic and fast fliers	insects (flies, moths, beetles, mosquitoes)		Grinnell 1962 Griffin 1958 Walker 1963
<i>Natalus mexicanus</i>	to 10 g	neotropical	Fluttering, mothlike flight	insects		Novick 1962 Walker 1968

I  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
Noctilio labialis minor	15-22 g	neotropical	Catch insects on water surface	insects (water beetles)		Griffin and Novick 1955 Walker 1968
Noctilio leporinus	30-70 g pinnae: 26 mm long 9 mm wide	streams, lakes, and ocean neotropical	Slow, deliberate flight. Gaffs fish by trailing feet through water	fish and insects		Grinnell 1962 Bloedel 1955 Griffin and Novick 1955 Griffin 1958 Walker 1968
Phyllostoma stenops	20-25 g	neotropical				Pye 1967 Walker 1968
Phyllostoma discolor	20-40 g	neotropical		fruit, nectar, pollen		Walker 1968
Phyllostoma hastatus panamensis	70-100 g pinnae: 28 mm long 12 mm wide	neotropical	Straight, rapid fliers	omnivorous (small animals-- rodents, birds, bats-- insects, fruit)		Grinnell 1970 Griffin 1958 Griffin and Novick 1955 Pye 1967 Grinnell 1962 Walker 1968

V  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>M. onyx vivax</i>		Gulf of California	(see remarks)	crustaceans and fish	Assumed bat gauffs fish by trailing feet in water	Bloedel 1955 Webster and Brazier 1968, 1969 Walker 1968
<i>Plecotus townsendii</i> ( <i>plecotus rafinesqii</i> )	10 g pinnae: 37 mm long 12 mm wide	temperate regions of New World	Versatile flyers; can hover in flight. Hunt along solid surfaces	insects (moths)		Grinnell 1962 Walker 1968
<i>Peronotus davyi</i>	7-10 g	near water neotropical	Swift flight near ground	insects		Novick 1965 Walker 1968
<i>Peronotus sumarensis</i>		feed along streams and rivers in wooded areas neotropical	Unusually skillful fliers; capable of hovering in flight	insects		Grinnell 1970 Walker 1968
<i>Rhynchiscus naso</i> ( <i>Rhynchonycteris naso</i> )	2-4 g	near water neotropical	Slow, steady flight just above water surface	insects		Griffin and Novick 1955 Griffin 1958 Walker 1968
<i>Saccopteryx bilineata</i>	5-6 g pinnae: 15 mm long 7.5 mm wide	neotropical	Fly in open spaces	insects (beetles and moths)		Griffin and Novick 1955 Grinnell 1962 Walker 1968

I  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Sturnira</i> sp.	16-20 g	forest (tropical and arid) neotropical		fruit		Pye 1967 Walker 1968
<i>Tadarida brasiliensis</i> <i>mexicana</i>	10-15 g pinnae: 9.5 mm long 7.5 mm wide	open country	High flyer	insects		Vernon and Peterson 1965 Grinnell 1962 Walker 1968
<i>Uroderma bilobatum</i>	13-21 g	neotropical		fruit		Walker 1968
<i>Vampyrus helleri</i>	12-15 g	watercourses in tropical forests neotropical		fruit		Walker 1968
<i>Vampyrus spectrum</i>	145-190 g	neotropical		small animals (birds, rodents, bats), insects, fruit		Walker 1968



### **APPENDIX III**

#### **TABULATIONS OF BIOLOGICAL SONAR DATA FOR WHALES, DOLPHINS, AND PORPOISES**

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	PEAK FREQUENCY kHz	BANDWIDTH <sup>(a)</sup> kHz	DURATION msec	SPL db re 1 $\mu$ bar	REMARKS	REFERENCES
<i>De. phinus delphis</i>	pulse (click)	30-60	to 150	0.05-0.25	40		Evans and Evans 1971 Evans 1972
<i>Echiorhynchus gibbosus</i>	pulse	2-3 kHz	to 120	1-1.5	72	*Upper limit of sonogram	Poulter 1968 Evans and Evans 1971
<i>Globicephala melana</i>	pulse (click)	5	~50	0.5		*10% threshold on Fourier power spectrum	Busnel et al. 1971
<i>Globicephala scammii</i>	pulse (click)	~3	to 160 to 100	0.5-2 0.25-2	80	*Upper limit of sonogram	Morris 1969 Evans and Evans 1971 Evans 1972
<i>Inia geoffrensis</i>	pulse (click)	60-75	21500	0.015-0.1	66	*10% threshold on Fourier power spectrum	AKL 1970
<i>Lagenorhynchus australis</i>	pulse (click)	~1 ~5-6	<2 to 160	1.5-5 0.8-1	~80	Limited hydrophone bandwidth (10-34); limited recorder bandwidth (30 kHz) *Upper limit of sonogram	Schevill and Watkins 1971
<i>Lagenorhynchus obliquidens</i>	pulse		to 80	0.25-1	70		Evans 1972
<i>Orcinus orca</i>	pulse (click)	~12	to 350	0.1-0.5	~80	*10% threshold on Fourier power spectrum	AKL 1971

<sup>(a)</sup>DEPARTMENTAL ANALYSIS

# I SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	PEAK FREQUENCY kHz	BANDWIDTH <sup>(1)</sup> kHz	DURATION msec	SPL dB re 1 $\mu$ bar	REMARKS	REFERENCES
Phocoena phocoena	pulse (click)	2	~4* to ~100 170-2160*	0.5-5 0.02-1.5 0.1	25-30 40	Limited bandwidth instrumentation: $\pm$ 10 kHz *Upper limit of sonagram Instrumentation capabilities unknown Two hydrophones: LC-32, MAX 8100 *Upper limit of spectrum analyzer trace	Schevill, Watkins, and Ray 1969 Dubrovski, Krasnov, and Titov 1971 *Phil and Andersen 1972
Phocoenoides dalli	pulse (click)	1-2	to 14*	0.5-1.5		Limited hydrophone bandwidth (LC-32) *Limit of trace on sonagram	Ridgway 1966 Evans and Evans 1971 Evans 1972
Physeter catodon	pulse (click)	~6	to 15*	0.75-5	to 84**	*Upper limit of sonagram *Limited bandwidth instrumentation **1/3 octave, 1 kHz band; system bandwidth: 4 kHz	Dunn 1969 Bachus and Schevill 1966 Evans 1972
Platanista gangetica	pulse (click)	20-40	to 150*	0.075-0.15	40	*Limit of trace on sonagram: 100 kHz	Herald et al. 1969 Evans 1972
Platanista indi	pulse (click)	50-60	to 100*	~0.02		Limited hydrophone bandwidth (LC-32): limited recorder bandwidth (30 kHz) *Upper frequency limit of instrumentation	Filleri, Kraus, and Gahr 1971
Pseudorca crassidens	pulse (click)		to 50*	0.2-0.5		*Upper frequency limit of instrumentation	Buanel and Dziedziec 1966

<sup>(1)</sup> DEFINITION: ANALYSIS

I  
SIGNAL CHARACTERISTICS

SPECIES	SIGNAL FORM	PEAK FREQUENCY kHz	BANDWIDTH <sup>(a)</sup> kHz	DURATION sec	SPL db re 1 $\mu$ bar	REMARKS	REFERENCES
<i>Stenella attenuata</i>	pulse		to 150	0.075-0.2			Evans and Evans 1971
<i>Stenella styx</i>	pulse		to 22			Limited bandwidth instrumentation	Buarel, Pilleri, and Frazer 1966
<i>Steno bredaensis</i>	pulse (click)	40-80	to 208 <sup>a</sup>	0.05-0.25		<sup>a</sup> Sonagram	Norris and Evans 1967 Evans and Evans 1971 Evans 1972
<i>Tursiops truncatus</i>	pulse (click)	30-60	to 130 <sup>a</sup>	0.02-0.1	75-110	<sup>a</sup> 10 <sup>4</sup> threshold on Fourier power spectrum	ARL/MUC 1969
<i>Tursiops truncatus</i> (Black Sea)	pulse (click)	~15		0.05-0.5		Instrumentation capabilities unknown	Airapetians et al. 1971

<sup>(a)</sup>DETERMINED BY ANALYSIS

# X SONAR CHARACTERISTICS

SPECIES	STANDARD SPEED, m/sec	PRR, /sec	BEAMWIDTH, deg	REMARKS	REFERENCES
Delphinus delphis	to 13		50-60 (60 kHz) (small only) ~75 (80 kHz) (small only) ~90 (100 kHz) (small only) ~15 (100 kHz) (small and melon)	Artificial source positioned at location of nasal plugs	Romanenko, Tomilin, and Artemenko 1965 Romanenko 1965 Walbert 1968
Inia geoffrensis	~1 4.5 (burst)	to 135		Swims laterally or vertically by linear motion with abrupt right angle turns	Fenner and Murchison 1970 ARL 1970 Layne 1958
Lagenorhynchus australis		to 25 (narrowband click) to 80 (broadband click)			Schervill and Watkins 1971
Orcinus vrea	to 15	>250	40		ARL 1971 Evans 1972 Johannessen and Harder 1960
Phocoena phocoena		to 1000	<60, (130 kHz)		Schervill, Watkins, and Ray 1969 Aphi and Andersen 1972
Phocoenoides dalli	>9	to 500		Click burst produced in captivity	Ridgway 1966

# X SONAR CHARACTERISTICS

SPECIES	SONAR SPEED, m/sec	PRR, /sec	BEAMWIDTH, deg	REMARKS	REFERENCES
<i>Physar catadon</i>	-2	to 50, average 3-7			Bachus and Schervill 1966 Dunn 1969 Walker 1968
<i>Platystia gnamptica</i>	-2-4	20-50		Serim on side with body axis inclined ~10° to bottom	Harold et al. 1969 Filleri 1970 Filleri, Kraus, and Gibb 1971
<i>Platoniata indl</i>		to 125			
<i>Pseudorca crassidens</i>		to 140			Bussell and Iniedsic 1966
<i>Stenella styx</i>		~60			Bussell, Filleri, and Frazier 1968
<i>Steno bredonensis</i>		to >100	20, (-5 db) ~4, (200 kHz) ~10, (150 kHz) ~60, (100 kHz)		Borris and Evans 1967 Evans 1967
<i>Tursiops truncatus</i>	-9	to 600	~25, (70 kHz) (horizontal: shall only) ~22, (70 kHz) (vertical: shall only) ~25-30 (horizontal: -3 db)	-3 db  0° tilt re rostrum axis	Evans, Sutherland, Bell 1964 Evans and Powell 1967 Evans 1972

# II ACOUSMETRIC DATA

SPECIES	AUSMOGRAM <sup>(1)</sup>	FREQUENCY RANGE, kHz	MAXIMUM SENSITIVITY			REMARKS	REFERENCES
			FREQUENCY, kHz	dB, dB in 0.001 sec/cm <sup>2</sup>	AREA		
Delphinus delphis	behavioral	to 280					Bel'kovich and Solntseva 1970
Inia geoffrensis	behavioral	1-105	75-90 <sup>a</sup> 30-50 <sup>a</sup>	24		<sup>a</sup> Turt; #Fig. 3 of reference 60 dB roll-off, 50 kHz to 100 kHz (Fig. 3 of reference).	Jacobs and Hall 1972
Orcinus orca	behavioral	0.5-51	15-25	0-10		60 dB roll-off, 25 kHz to 51 kHz	Hall and Johnson 1972
Phocoena phocaena	behavioral	1-150	8-32	~24		15 dB/octave roll-off, 40 kHz to 140 kHz; 700 dB/octave roll-off, 140 kHz-150 kHz	Andersen 1970
Stenella attenuata	electrophysiological (E <sub>h</sub> )	10-150	50-70	5-20	lower jaw	70 dB roll-off, 70 kHz to 140 kHz	Bullock et al. 1968
Stenella caeruleoalba	electrophysiological (E <sub>h</sub> )	10-150	50-70	5-20	lower jaw	70 dB roll-off, 70 kHz to 140 kHz	Bullock et al. 1968
Tursiops gilli	electrophysiological (E <sub>h</sub> )			10-25	lower jaw	Dependent upon stimulus duration	Bullock et al. 1968 McDermick 1968 McDermick et al. 1970 Morris 1969
Tursiops truncatus	behavioral	0.1-150	30-80	~10		Dependent upon stimulus duration	Johnson 1966, 1968

<sup>(1)</sup> BEHAVIORAL: TALK  
ELECTROPHYSIOLOGICAL

**II**  
**DETECTION/DISCRIMINATION DATA**

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION				SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION			REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	SIGNAL FORM		
Delphinus delphis	3-step pyramids, styrofoam	base: 100 cm <sup>2</sup> each; 2nd step 49 cm <sup>2</sup> each; 3rd step 9 cm <sup>2</sup> target 1 8.6 cm <sup>2</sup> target 2 stage 1.2 thick	discrimina- tion, size (6.7% diff.)	600-800	>50%			*Value not reported	Gurevich 1969
	squares, styrofoam	100 cm <sup>2</sup> versus 90.65 cm <sup>2</sup> ; 1.2 thick	discrimina- tion, area (9.75% diff.)	600-800	>50%				
	3-step pyramids, styrofoam ebonite	same as target 1 above	discrimina- tion, material		~100%				
	squares, styrofoam ebonite	100 cm <sup>2</sup>	discrimina- tion, material		~100%				
fish (bullhead) versus tube, rubber	fish (bullhead)	12-17	discrimina- tion	150-500	85% (60 trials)				Konstantinov, Mel'nikov, and Titov 1968
	fish (bullhead)	4, diam 22, long 12-17 versus 27-30	discrimina- tion, size	260, average	88%			*Smaller fish positive target (edible)	

<sup>(1)</sup>BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL



IV  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION				SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	SIGNAL FORM	TASK	PERFORMANCE	
Delphinus delphis (Cont'd)	fish (bullhead) versus fish (metal clad)	equal size	discrimina- tion, material	300, average	~85%					
	square versus triangle, ebonite and plexiglass	100 cm <sup>2</sup> 50 cm <sup>2</sup>	discrimina- tion, size and shape	800	100%					Magdonas Bel'kovich and Krushinskaya 1970
Inia geoffrensis	wire, metal	0.26 diam 0.13 0.12 0.11	detection	219	100% 73% 63% ~45%					Permer and Murchison 1970
	cylinder, copper	7.6 long 4.2 diam versus 5.6 diam	discrimina- tion, size*		75%				*2 dB target strength difference	Evans 1972
Lagenorhynchus obliquatus	ring, plastic (hollow)	25 diam	detection	200						Evans 1972
	discs, copper versus aluminum	50 diam	discrimina- tion, material		>50%				*Value not reported	Evans 1972
Orca orca	ring, plastic (hollow)	25 diam	detection	200						Evans 1972

<sup>(1)</sup> DENAVISAL: TASK  
ELECTROPHYSIOLOGICAL

# IV DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE; DETECTION/DISCRIMINATION				SIGNAL; DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	SIGNAL FORM	TASK	PERFORMANCE	
Phocoena phocoena	wire, metal (barrier, 1 m separation)	0.05 diam 0.075 diam	avoidance		≥90% 79% ~50%	behavioral	tone burst, 2 kHz	detection, angular displacement of signal (NMA)	75%, 5°, median plane	Bunnell, Dziedziec, and Andersen 1965 Bunnell and Dziedziec 1967 Andersen 1970
	lime, noncellulose nylon (barrier, 1 m separation)	0.18 diam 0.15 diam	avoidance		85% 72%					
	lime, 3-strand perlon (barrier, 1 m separation)	0.1 diam 0.08 diam	avoidance		>95% 58%					
Turriopsis truncatus (Black Sea)	cylinder	11 diam 25 long versus 28	discrimination, length		100%, 25 versus 235 70%, 25 versus 30 50%, 25 versus 28	behavioral		discrimination, target range discrimination, angular separation of targets	Δθ=1.5 mm Δφ=0.25° (horizontal) 0.8° (vertical)	Ayrapet'yants et al. 1969 Akrapet'yants and Konstantinov 1971

<sup>(1)</sup> BEHAVIORAL; TASK  
ELECTROPHYSIOLOGICAL

IV  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE: DETECTION/DISCRIMINATION				SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION			REMARKS	REFERENCES
	FORM/MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(1)</sup>	TASK	PERFORMANCE	
Tursiops truncatus (Montagu)	spheres, steel	5.4 diam 5.71 diam 6.35 diam	discrimination, size	45-60	100%, 6.35 versus 5.4 77%, 6.35 versus 5.71				Norris, Evans, and Turner 1967
	discs, metal: copper	30 diam thick- ness: 0.22 std 0.32 0.27 0.16 0.64 0.32	discrimination, material, thickness		75%, 0.22 versus 0.32 60%, 0.22 versus 0.27 50%, 0.22 versus 0.16				Evans and Powell 1967
	brass				100%, 0.22 versus 0.64 55%, 0.22 versus 0.32				
	aluminum	0.79 0.64 0.32			99%, 0.22 versus 0.79 97%, 0.22 versus 0.64 100%, 0.22 versus 0.32				
	cylinder, vitamin capsule	0.5 diam 2.5 long	Detection	~50-60					Johnson 1967
	cylinders, corprene	17.8 long 4.2 diam versus 5.2 diam	discrimination, size <sup>a</sup>		85%			Target lying on tank floor	
	gelatin capsule, water-filled with pebbles versus fish muscle	3.8 long	discrimination, material	~50-50	100%			*1 dB target strength difference	Evans 1972
									Norris et al. 1961

<sup>(1)</sup> BEHAVIORAL TASK  
ELECTROPHYSIOLOGICAL

IV  
DETECTION/DISCRIMINATION DATA

SPECIES	TARGET/OBSTACLE: DETECTION/DISCRIMINATION				SIGNAL: DETECTION, DISCRIMINATION, LOCALIZATION				REMARKS	REFERENCES
	FORM/ MATERIAL	SIZE, cm	TASK	DISTANCE, cm	PERFORMANCE	TECHNIQUE <sup>(a)</sup>	SIGNAL FORM	TASK	PERFORMANCE	
Tursiops truncatus (Montagu) (Cont'd)	discs, cellite	diff diam, 15.2 std	discrimina- tion, size	120-180	58%, $\Delta$ diam 0.5 cm <sup>2</sup> 75%, $\Delta$ diam 0.9 cm <sup>2</sup>	behavioral	tone burst	discrimina- tion, intensity	$\Delta I$ -1 dB	* $\Delta$ area: 6.7% and 12% $\Delta$ target strength: 0.6 dB and 1 dB
	disc versus square versus triangle, cellite	15.2 diam equal area	discrimina- tion, shape	120-180	>95%					
	fish	5-15 long	detection	500-1000						Evans 1972

<sup>(a)</sup> BEHAVIORAL: TASK  
ELECTRON: BIOLOGICAL

I  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Delphinus delphis</i>	to 75 kg	worldwide pelagic	Very fast swimmer. Sons in rectilinear motion--up-down or sideways	fish and cephalopods found in shoals and near surface	Travel in groups from ~20 to several hundred	Walker 1968 Konstantinov, Mel'nikov, and Titov 1968
<i>Globicephala</i> sp.		offshore to inshore waters (seasonal) pelagic to littoral		squid, fish, cuttlefish	Schools may contain several hundred animals. During feeding schools may disperse	Walker 1968 Leatherwood, Evans, and Rice 1972 Morris 1969
<i>Inia geoffrensis</i>	to 125 kg	Amazon and Orinoco Rivers riverine	Sons in rectilinear motion--up-down or sideways. Hunts at leisurely swimming speed (2-3 kt)	fish (<30 cm)	Feed in small groups (3-6 animals). Habitat is muddy with minimal visibility	Walker 1968 Penner and Marchison 1970 Layne 1958
<i>Leptorhynchus</i> sp.	to 150 kg	offshore to inshore waters (seasonal) pelagic to littoral	Are vigorous swimmers	fish--herring, mackerel, anchovies-- crustaceans, squid	Some species feed in large schools (1000-2000 animals)	Walker 1968 Leatherwood, Evans, and Rice 1972
<i>Orcinus orca</i>	1000-2000 kg	worldwide, principally Arctic and Antarctic pelagic to estuarine, sometimes riverine	Apparently sons by rolling from side to side	small marine mammals (seal, walrus, porpoise), baleen whales, birds, fish, cephalopods		Walker 1968 Evans 1972 Hall and Johnson 1972 Schervill and Watkins 1966

I  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
<i>Phocoena phocoena</i>	50-75 kg	European and African waters. Littoral and sometimes riverine	Usually swims just below surface. Feeds near bottom	fish (<30 cm), cephalopods, crustaceans	*Have been caught in fishnets at depths of 75 m	Walker 1968 Dubrovskii, Krasnov, and Titov 1971
<i>Phocoenoides dalli</i>	80-125 kg	pelagic	Very fast swimmer	fish and squid	Found in small groups (2-20 animals)	Walker 1968 Ridgway 1966
<i>Physeter catodon</i>	30-45x10 <sup>3</sup> kg	pelagic	Feed at great depths (~500 m)	squid, cuttlefish, fish (barracuda, albacore), sharks	Swim (hunt) in groups of 15-20 animals	Walker 1968
<i>Platanista gangetica</i>	20-225 kg	Ganges, Indus, and Brahmaputra Rivers riverine	Swim on side with flipper sometimes dragging along bottom. Average swimming speed ~3 kt	fish (<30 cm) and crustaceans (shrimp)	Habitat is muddy with minimal visibility. Hunt singly or in small groups (2-10 animals)	Walker 1968 Merald et al. 1969 Filleri 1970
<i>Platanista indi</i>	20-225 kg	Ganges, Indus, and Brahmaputra Rivers riverine	Swim on side with flipper sometimes dragging along bottom. Average swimming speed ~3 kt	fish (<30 cm) and crustaceans (shrimp)	Habitat is muddy with minimal visibility. Hunt singly or in small groups (2-10 animals)	Walker 1968 Merald et al. 1969 Filleri 1970
<i>Pseudorca crassidens</i>	1400 kg	pelagic		cephalopods and fish	Feed in groups of several hundred animals	Walker 1968

Y  
HABITAT/PREY/BIOLOGY

SPECIES	SIZE	HABITAT	BEHAVIOR	PREY	REMARKS	REFERENCES
Stenella sp.	to 165 kg	pelagic	Hunt (feed) near surfaces	fish	Occur in schools of a few to several hundred animals	Walker 1968 Leatherwood, Evans, Rice 1972
Tursiops truncatus (Moutagu)	150-200 kg	bays and inshore waters littoral and sometimes riverine	Tilts head downward and scans laterally over 25-50° arc or with a circular motion while echolocating	fish (most abundant species), sharks, shrimp, rays, squid		Morris et al. 1961 Evans and Prescott 1962 Walker 1968

#### APPENDIX IV

#### BIBLIOGRAPHY OF ACCUMULATED LITERATURE FOR BATS



LISTING FORMAT

		LANGUAGE	
6-DIGIT CODE		ORIGINAL	TRANSLATION
AUTHOR(S)	YEAR OF ISSUE		
TITLE OR DESCRIPTION			
SOURCE			

Preceding page blank

# KEY TO 6-DIGIT NUMBER CODE

## DIGITS

1	2	3	4	5	6
---	---	---	---	---	---

## DIGIT(S)

1

## FUNCTION(S)

### ORDER

- 1 CETACEA
- 2 CHIROPTERA
- 4 OTHER

2-3

### ECHOLOCATION

- 1 DETECTION
- 2 DISCRIMINATION
- 10 PERFORMANCE
- 20 SIGNALS ; FORMAT
- 40 BEHAVIOR

4-5

### BIOLOGY

- 1 MORPHOLOGY ; ANATOMY
- 2 AUDIOMETRY
- 4 SIGNAL PROCESSING, CONDITIONING
- 10 SOUND GENERATION
- 20 SOUND RECEPTION

6

### ECOLOGY

- 1 HABITAT
- 2 PREY

**Examples:**

- 231210** A report about bats (2). Includes measurements, description, or commentary about echolocation signals (e.g., characteristics, repetition rate) (20), and the bats' performance (10) on target or obstacle detection or avoidance (1) ( $20+10+1=31$ ), plus measurements, description or commentary on the bats' ability or capability to detect and/or localize sounds (echoes) (20), and the morphology and/or anatomy of the hearing organ (1) ( $20+1=21$ ). Does not include information about the bats' normal hunting habitat (e.g., food prey) (0).
- 572101** A report about whales, dolphins, or porpoises (1), including measurements and/or commentary on other animal forms (4) ( $4+1=5$ ). Includes measurements, description, or commentary about the animals' echolocation signals (20), and their performance(s) (10) in discriminating between targets (2), plus commentary on the animals' behavior while echolocating (e.g., scanning motions, swimming speed) (40) ( $40+20+10+2=72$ ). Includes commentary on the mechanics of sound generation (e.g., beaming, energy requirements) (10), plus description of the animals' normal hunting habitat(s) (1).

- 203240 ENG  
AIDLEY, U. J. 1959  
[ECHO INTENSITY IN RANGE ESTIMATION BY BATS]  
NATURE, 224, 1330-1331
- 263260 RUS ENG  
AIRAPETIANZ, E. SH., 1967  
BIONIC ASPECTS OF RESEARCH INTO MECHANISMS RESPONSIBLE FOR SPATIAL  
INFORMATION ECHO-LOCATION IN BATS]  
FIZIOLOGICHESKII ZH. SSSR MOSCOW, 54, 368-376
- 250000 RUS ENG  
AIRAPETIANZ, E. SH., AND A. I. KONSTANTINOV 1965  
[ON THE QUESTION OF THE ROLE OF ECHOLOCATION IN SPATIAL ANALYSIS  
BY BATS]  
BIONIKA, NAUKA PUBL. HOUSE, MOSCOW, 334-341
- 263000 RUS ENG  
AIRAPETIANZ, E. SH., A. I. KONSTANTINOV 1967  
[ON THE INTERACTION OF ANALYZERS IN THE ECHOLOCATION ACTIVITIES  
OF BATS]  
RESEARCH IN BRAIN SIGNALIZATION APPARATUS. NAUKA, Leningrad, 22-32
- 220340 RUS ENG  
AIRAPETIANZ, E. SH., A. I. KONSTANTINOV 1968  
[RESEARCH ON THE NERVOUS ECHOLOCATION MECHANISM IN BATS]  
VOPROSY BIONIKI [PROBLEMS OF BIONICS] 445-453
- 273250 ENG  
AIRAPETIANZ, E. SH., A. I. KONSTANTINOV, D. P. MATJUSHKIN 1969  
[BRAIN ECHOLOCATION MECHANISMS AND BIONICS]  
ACTA PHYSIOL. ACAD. SCI. HUNG., 35, 1-17
- 313250 ENG  
AIRAPETIANZ, E. SH., A. I. KONSTANTINOV 1971  
[PHYSIOLOGICAL INVESTIGATIONS OF ULTRASONIC ECHOLOCATION IN  
ANIMALS]  
PAPER PRESENTED AT THE 25TH INTERNATL. CONG. OF PHYSIOL. SCIENCES,  
MUNICH, JULY 1971
- 273230 ENG  
AIRAPETIANZ, E. SH., A. I. KONSTANTINOV 1968  
[ECHOLOCATING SPACE ANALYSIS OF ANIMALS]  
VOLUNTEER ABSTRACTS, 8, 5 (24TH INTERNATL. CONG. PHYSIOL. SCI.,  
WASH., D.C.)
- 222000 ENG  
ALTES, R. A. AND E. L. TITLERAUM  
[HAT SIGNALS AS OPTIMALLY DOPPLER TOLERANT WAVEFORMS]  
J. ACOUST. SOC. AM. 48, 1014-1017



271000 ENG  
CHASE, J., R.A. SUTHERS 1969  
[VISUAL OBSTACLE AVOIDANCE BY ECHOLOCATING BATS]  
ANIMAL BEHAVIOR 17, 201-207

273000 ENG  
CURTIS, W.E. 1952  
[QUANTITATIVE STUDIES OF ECHOLOCATION IN BATS (MYOTIS LUCIFUGUS),  
STUDIES OF VISION OF BATS (MYOTIS LUCIFUGUS AND EPTESICUS F.  
FUSCUS) AND QUANTITATIVE STUDIES OF VISION OF OWLS (TYTO ALBA  
PRATINCOLA)]  
THESIS PRESENTED TO FACULTY OF GRAD. SCHOOL OF CORNELL UNIV.  
200220 ENG  
DALLAND, J.I. 1965  
[AUDITORY THRESHOLDS IN THE BAT-A BEHAVIORIAL TECHNIQUE]  
J. AUD. RES. 5, 95-108

200220 ENG  
DALLAND, J.I. 1965  
[HEARING SENSITIVITY IN BATS]  
SCIENCE 150, 1185-1186

200220 ENG  
DALLAND, J.I., J.A. VERNON, E.A. PETERSON 1966  
[HEARING AND COCHLEAR MICROPHONIC POTENTIALS IN THE BAT EPTESICUS  
FUSCUS]  
J. NEUROPHYSIOL. 30, 697-709

271003 GER ENG  
DIJKGRAAF, V.I. 1946  
[DIE SINNESWELT DER FLEDERMAUSE (THE SENSORY WORLD OF RATS)]  
EXPERIENTIA 2, 438-448

200002 ENG  
DUNN, L.  
[OBSERVATIONS ON THE CARNIVOROUS HABITS OF THE SPEAR NOSED BAT,  
PHYLLOSTOMUS HASTATUS PANAMENSIS ALLEN, IN PANAMA]  
J. MAMM. 14, 188-195

241000 ENG  
GALAMBOS, R. 1942  
[THE AVOIDANCE OF OBSTACLES BY FLYING BATS- SPALLANZANIS IDEAS  
[1794] AND LATER THEORIES]  
ISIS 34, 132-14

271110 ENG  
GALAMBOS, R., D. R. GRIFFIN 1942  
[OBSTACLE AVOIDANCE BY FLYING BATS, THE CRIES OF BATS]  
J. EXP. ZOOL. 84, 475-490

261002	GOULD, E. 1955 [THE FEEDING EFFICIENCY OF INSECTIVOROUS BATS] J. MAMM. 36, 399-407	ENG
240000	GOULD, E. 1959 [FURTHER STUDIES ON THE FEEDING EFFICIENCY OF BATS] J. MAMM. 40, 149-150	ENG
220110	GRIFFIN, D.R. 1946 [SUPERSONIC CRIES OF BATS] NATURE 158, 46-48	ENG
220000	GRIFFIN, D.R. 1950 [MEASUREMENTS OF THE ULTRASONIC CRIES OF BATS] J. ACOUST. SOC. AM. 22, 247-255	ENG
230000	GRIFFIN, D.R. 1950 [THE NAVIGATION OF BATS] SCI. AM. 183, 52-55	ENG
220000	GRIFFIN, D.R. 1951 [AUDIBLE AND ULTRASONIC SOUNDS OF BATS] EXPERIENTIA 7, 448-453	ENG
220110	GRIFFIN, D.R. 1952 [MECHANISMS IN THE BAT LARYNX FOR PRODUCTION OF ULTRASONIC SOUNDS] PROC. AM. PHYSIOL. SOC. [ABSTRACT] 11, 56	ENG
263001	GRIFFIN, D.R. 1953 [BAT SOUNDS UNDER NATURAL CONDITIONS, WITH EVIDENCE FOR ECHOLOCATION OF INSECT PREY] J. EXP. ZOOL. 123, 435-466	ENG
273373	GRIFFIN, D.R. 1958 [LISTENING IN THE DARK] YALE UNIV. PRESS, NEW HAVEN	ENG
273310	GRIFFIN, D.R. 1959 [ECHOES OF HATS AND MEN] ANCHOR BOOKS, DOUBLEDAY AND CO., INC., GARDEN CITY, NEW YORK	ENG

- 221000 ENG  
GRIFFIN, D.R. 1962  
[COMPARATIVE STUDIES OF THE ORIENTATION SOUNDS OF BATS]  
SYMP. ZOOL. SOC. LOND. 7, 61-72
- 252000 ENG  
GRIFFIN, D.R. 1967  
[DISCRIMINATIVE ECHOLOCATION BY BATS]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS VOL. 1, R.G. BUSNEL [ED.],  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 273-305
- 221001 ENG  
GRIFFIN, D.R. 1971  
[THE IMPORTANCE OF ATMOSPHERIC ATTENUATION FOR THE ECHOLOCATION  
OF BATS [CHIROPTERA]]  
ANIMAL BEHAVIOUR 19, 55-61
- 251000 ENG  
GRIFFIN, D.R., R. GALAMBOS 1941  
[THE SENSORY BASES OF OBSTACLE AVOIDANCE BY FLYING BATS]  
J. EXP. ZOOL. 86, 481-506
- 261013 ENG  
GRIFFIN, D.R., A. NOVICK 1955  
[ACOUSTIC ORIENTATION OF NEOTROPICAL BATS]  
J. EXP. ZOOL. 130, 251-300
- 211000 ENG  
GRIFFIN, D.R., A.D. GRINNELL 1958  
[ABILITY OF BATS TO DISCRIMINATE ECHOES FROM LOUDER NOISES]  
SCIENCE 128, 145-46
- 271100 ENG  
GRIFFIN, D.R., A. NOVICK, M. KORNFIELD 1958  
[THE SENSITIVITY OF ECHOLOCATION IN THE FRUIT BAT, ROUSETTUS]  
BIOL. BULL. 115, 107-113
- 271002 ENG  
GRIFFIN, D.R., F.A. WEBSTER, C.R. MICHAEL 1960  
[THE ECHOLOCATION OF FLYING INSECTS BY BATS]  
ANIMAL BEHAVIOUR, 8, 141-154
- 260210 ENG  
GRIFFIN, D.R., D.C. DUNNING, A.D. CAHLANDER, F.A. WEBSTER, 1962  
[CORRELATED ORIENTATION SOUNDS AND E.R. MOVEMENTS OF HORSESHOE BATS]  
NATURE 196, 1185-1188
- 271230 ENG  
GRIFFIN, D.R., J.J.G. MC CUE, A.D. GRINNELL 1963  
[THE RESISTANCE OF BATS TO JAMMING]  
J. EXP. ZOOL. 152, 229-250



- 273002 ENG  
GRIFFIN, D.R., J.H. FRIEND, F. WEBSTER 1965  
[TARGET DISCRIMINATION BY THE ECHOLOCATION OF BATS]  
J. EXP. ZOOL. 158, 155-168
- 233270 ENG  
GRINNELL, A.D. 1962  
[NEUROPHYSIOLOGICAL CORRELATES OF ECHOLOCATION IN BATS]  
HARVARD BIBLIOGRAPHICAL LABS TR 30, PROJ. NR-301-219, CONT. NONR-1866  
[12] PP. 123, AD 686 044
- 200220 ENG  
GRINNELL, A.D. 1963  
[THE NEUROPHYSIOLOGY OF AUDITION IN BATS, INTENSITY AND FREQUENCY  
PARAMETERS]  
J. PHYSIOL. 167, 38-66
- 200240 ENG  
GRINNELL, A.D. 1963  
[THE NEUROPHYSIOLOGY OF AUDITION IN BATS, TEMPORAL PARAMETERS]  
J. PHYSIOL. 167, 67-96
- 200240 ENG  
GRINNELL, A.D. 1963  
[THE NEUROPHYSIOLOGY OF AUDITION IN BATS, DIRECTIONAL LOCALIZATION  
AND BINAURAL INTERACTION]  
J. PHYSIOL. 167, 97-113
- 203260 ENG  
GRINNELL, A.D. 1967  
[MECHANISMS OF OVERCOMING INTERFERENCE IN ECHOLOCATING ANIMALS]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL (ED.),  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 451-478
- 220220 ENG  
GRINNELL, A.D. 1970  
[COMPARATIVE AUDITORY NEUROPHYSIOLOGY OF NEOTROPICAL BATS  
EMPLOYING DIFFERENT ECHOLOCATION SIGNALS]  
Z. VERGL. PHYSIOL. 68, 117-153
- 231000 ENG  
GRINNELL, A.D., D.R. GRIFFIN 1958  
[THE SENSITIVITY OF ECHOLOCATION IN BATS]  
BIOL. BULL. 114, 10-22
- 200230 ENG  
GRINNELL, A.D., J.J.G. MC CUE 1963  
[NEUROPHYSIOLOGICAL INVESTIGATIONS OF THE RAT, MYOTIS LUCIFUGUS,  
STIMULATED BY FREQUENCY MODULATED ACOUSTICAL PULSE]  
NATURE 194, 453-455

- 200210  
ENG  
GRINNELL, A.D., V.S. GRINNELL 1965  
[NEURAL CORRELATES OF VERTICAL LOCALIZATION BY ECHOLocATING BATS]  
J. PHYSIOL. 181, 830-851
- 220250  
ENG  
GRINNELL, A.D., S. HAGIWARA 1972  
[ADAPTATIONS OF THE AUDITORY NERVOUS SYSTEM FOR ECHOLocATION]  
Z. VERGL. PHYSIOL. 76, 41-81
- 220220  
ENG  
GRINNELL, A.D., S. HAGIWARA 1972  
[STUDIES OF AUDITORY NEUROPHYSIOLOGY IN NON-ECHOLocATING BATS AND  
ADAPTATIONS FOR ECHOLocATION IN ONE GENUS, ROUSSETTUS]  
Z. VERGL. PHYSIOL. 76, 82-96
- 251010  
ENG  
GRUMMON, R.A., A. NOVICK 1963  
[OBSTACLE AVOIDANCE IN THE BAT, MACROTUS MEXICANUS]  
PHYSIOL. ZOOL. 36, 361-369
- 251000  
ENG  
HAHN, W.L. 1908  
[SOME HABITS AND SENSORY ADAPTATIONS OF CAVE INHABITING BATS, I + II  
BIOL. BULL. 15, 135-193
- 200210  
ENG  
HENSON, O.W., JR. 1961  
[SOME MORPHOLOGICAL AND FUNCTIONAL ASPECTS OF CERTAIN STRUCTURES  
OF THE MIDDLE EAR IN BATS AND INSECTIVORES]  
UNIV. OF KANSAS SCIENCE BULL. 42, 151-255
- 200240  
ENG  
HENSON, O.W., JR. 1965  
[THE ACTIVITY AND FUNCTION OF THE MIDDLE-EAR MUSCLES IN ECHO-  
LOCATING BATS]  
J. PHYSIOL. 180, 871-887
- 200240  
ENG  
HENSON, O.W., JR. 1966  
[COMPARATIVE PHYSIOLOGY OF MIDDLE EAR MUSCLE ACTIVITY DURING  
ECHOLocATION IN BATS]  
REPRINTED FROM AMER. ZOOLOGIST [ABSTRACT] 6
- 221270  
ENG  
HENSON, O.W., JR. 1967  
[THE PERCEPTION AND ANALYSIS OF BIOSONAR SIGNALS BY BATS]  
ANIMAL SONAR SYSTEMS- BIOLOGY AND BIONICS, VOL.2, R.G. BUSNEL [ED.]  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 949-1003

- 200220 ENG  
HENSON, O.W. JR. 1967  
[AUDITORY SENSITIVITY IN MOLESTIDAE [CHIROPTERA]]  
ANAT. REC. [ABSTRACT] 157, 363-364
- 200270 ENG  
HENSON, O.W. JR. 1970  
[EAR AND AUDITION]  
CH. 4 IN THE BIOLOGY OF BATS, VOL. 2, W.A. WIMSATT [ED.]  
ACADEMIC PRESS, NEW YORK, 181-263
- 201000 ENG  
KAY, L., J.D. PYE, J. NORDMARK 1961  
[THE PERCEPTION OF DISTANCE IN ANIMAL ECHO-LOCATION]  
NATURE 190, 361-364
- 222000 ENG  
KAY, L. 1962  
[A PLAUSIBLE EXPLANATION OF THE BATS ECHO-LOCATION ACUITY]  
ANIMAL BEHAVIOUR 10, 34-41
- 220000 ENG  
KAY, L., T.J. PICKVANCE 1963  
[ULTRASONIC EMISSIONS OF THE LESSER HORSESHOE BAT RHINOLOPHUS  
HIPPOSIDEROS [RECH.]]  
PROC. ZOOL. SOC. LOND. 141, 163-171
- 250000 RUS ENG  
KONSTANTINOV, A.I. 1964  
[INFLUENCE OF BLOCKING VISUAL RECEPTION ON THE DIRECTED BEHAVIOR  
VESTNIK Leningrad Univ. 15, 72-75]
- 250010 RUS ENG  
KONSTANTINOV, A.I. 1965  
[THE INFLUENCE OF THE PARTIAL AND TOTAL ELIMINATION OF THE BRAIN  
CORTEX ON BAT ECHOLOCATION]  
DAN SSSR, 161, 989-991
- 253310 RUS ENG  
KONSTANTINOV, A.I. 1965  
[MATERIALS ON THE PHYSIOLOGY OF SPATIAL ANALYSIS IN BATS]  
AVTOREF. KAND. DISS. IZD-VO AN SSSR. PP. 17
- 271000 RUS ENG  
KONSTANTINOV, A.I. 1965  
[PRINCIPLES OF ULTRASOUND SPATIAL ORIENTATION IN BATS]  
VOPR. SRVNIIT. FIZIOL. ANALIZATOROV. [PROBLEMS OF THE COMPARATIVE  
PHYSIOL. OF ANALYZERS] LENINGRAD, 11, 93-111]

- 261002 RUS ENG  
KONSTANTINOV, A.I., 1969  
[RELATIONSHIP BETWEEN AUDITORY PERCEPTION AND ECHOLOCAION DURING  
HUNTING IN MYOTIS OXYGNATHUS]  
ZH. EVOL. BIOK. I FIZIOL. 5, 566-572
- 211200 RUS ENG  
KONSTANTINOV, A.I., B.V. SOKOLOV, I.M. STOSMAN 1967  
[COMPARATIVE RESEARCH ON ECHOLOCAION SENSITIVITY IN BATS]  
DAN SSSR 175, 1418-1421
- 253000 RUS ENG  
KONSTANTINOV, A.I., N.I. AKHMAROVA 1968  
[TARGET DISCRIMINATION BY ECHOLOCAION OF BATS]  
NAUC. DOKL. VYS. SHK. BIOL. NAUK. 11, 22-28
- 220100 RUS ENG  
KONSTANTINOV, A.I., B.V. SOKOLOV 1969  
[CHARACTERISTICS OF ULTRASONIC ORIENTATION SIGNALS IN HORSESHOE  
BATS [RHINOLOPHIDAE]]  
ZH. EVOL. BIOK. I FIZIOL. 5, 90-97
- 200220 RUS ENG  
KONSTANTINOV, A.I., N.N. SANOTSKAYA, N.N. SOKOLOVA  
FREQUENCY-THRESHOLD CHARACTERISTICS OF THE AUDITORY SYSTEM OF BATS  
J. HIGHER NERV. ACT. 21, 535-541
- 200200 RUS ENG  
KRAVTSOV, B.G., B.M. ZVONOV 1969  
[THE DEPENDENCE OF THE SUMMATED REACTION OF BAT AUDITORY NERVE ON  
THE SPATIAL POSITION OF THE SOUND SOURCE]  
MOSC. UNIV. VESTN. SER. 6, BIOLOGIIA POCH., 99-101
- 200110 GER ENG  
KULZER, E. 1956  
[FLYING FOX PRODUCES ORIENTATION SOUNDS WITH TONGUE]  
DIE NATURWISS. 43, 117-118
- 200030 ENG  
MC CUE, J.J. 1966  
[AURAL PULSE COMPRESSION BY BATS AND HUMANS]  
J. ACOUST. SOC. AM. 40, 545-548
- 200040 ENG  
MC CUE, J.J. 1969  
[SIGNAL PROCESSING BY THE BAT]  
J. AUD. RES. 9, 100-107

223240

FRE ENG

MERMOZ, H 1967  
[DISCUSSION OF PAPER BY A. GRINNELL]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUISNEL [ED.],  
LAB. DE PHYSIOL. ACOUST. JOUY-EN-JOSAS, FRANCE, 482-490  
233210 ENG

MOGUS, M.A. 1967  
[THEORIES OF BAT ECHOLOCACTION]  
ORDNANCE RESEARCH LAB., PENNA. STATE UNIV., TECH. MEMO. 657.2341-02,  
PP. 62 (15 FEBRUARY 1967) AD 650 476  
201230 ENG

MOGUS, M.A. 1970  
[A THEORETICAL APPROACH TO BAT ECHOLOCACTION]  
PENNA. STATE UNIV., DISSERTATION, UNIV. MICROFILMS, ANN ARBOR, MICH.  
241000 GER ENG

MOHRES, F.P. 1950  
[ZUR ORIENTIERUNG DER FLEDERMAUSE]  
NATUR UND VOLK, 80, 153-161  
261310 GER ENG

MOHRES, F.P. 1951  
[UBER EINE NEUE ART VON ULTRASCHALL-ORIENTIERUNG BEI  
FLEDERMAUSEN]  
VERH. DTSCH. ZOOL. 179-186  
261110 GER ENG

MOHRES, F.P. 1952  
[DIE ULTRASCHALL-ORIENTIERUNG DER FLEDERMAUSE]  
DIE NATURWISS. 39, 273-279  
240100 GER ENG

MOHRES, F.P. 1953  
[ULTRASCHALLORIENTIERUNG AUCH BEI FLUGHUNDEN [MACROCHIROPTERA-  
PTEROPODIDAE]] [ULTRASONIC ORIENTATION IN FLYING FOX BATS]  
DIE NATURWISS. 40, 536-537  
260310 GER ENG

MOEHRES, F.P. 1953  
[UBER DIE ULTRASCHALLORIENTIERUNG DER HUFSENNASEN [CHIROPTERA-  
RHINOLOPHINAE]]  
Z. VERGL. PHYSIOL. 34, 547-588  
260310 ENG

MOEHRES, F.P. 1960  
[SONIC ORIENTATION OF BATS AND OTHER ANIMALS]  
SYMP. ZOOL. SOC. LOND. 3, 57-66

- 231012  
 MOHRES, F.P. 1947  
 [ULTRASONIC ORIENTATION IN MEGADERMATID BATS]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
 LAB. DE PHYSIOL. ACOUST. JOUY-EN-JOSAS, FRANCE, 115-126  
 ENG
- 221000  
 MOHRES, F.P. 1947  
 [GENERAL CHARACTERS OF ACOUSTIC ORIENTATION SIGNALS AND  
 PERFORMANCE OF SONAR IN THE ORDER CHIROPTERA]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
 LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 401-407  
 ENG
- 262000  
 MOHRES, F.P. 1947  
 [COMMUNICATIVE CHARACTER OF SONAR SIGNALS IN BATS]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 2, R.G. BUSNEL [ED.],  
 LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 939-944  
 GER ENG
- 271010  
 MOHRES, F.P., G. NEUWEILER 1966  
 [DIE ULTRASCHALLORIENTIERUNG DER GROSSBLATT-FLEDERMAUSE  
 [CHIROPTERA-MEGADERMATIDAE]]  
 Z. VERGL. PHYSIOL. 53, 195-227  
 ENG
- 651101  
 NEUWEILER, G. 1967  
 [INTERACTION OF OTHER SENSORY SYSTEMS WITH THE SONAR SYSTEM]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1,  
 R.G. BUSNEL [ED.], LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE,  
 509-529  
 GER ENG
- 200220  
 NEUWEILER, G. 1970  
 [NEUROPHYSIOLOGISCHE UNTERSUCHUNGEN ZUM ECHOORTUNGSSYSTEM DER  
 GROSSEN HUFEISENNASE RHINOLOPHUS FERRUM EQUINUM SCHREIBER, 1774]  
 Z. VERGL. PHYSIOL. 67, 273-306  
 ENG
- 200250  
 NEUWEILER, G. 1970  
 [NEUROPHYSIOLOGICAL INVESTIGATIONS IN THE COLLICULUS INFERIOR OF  
 RHINOLOPHUS FERRUM EQUINUM]  
 BIJDRAAGEN TOT DE DIERKUNDE 40, 59-61  
 ENG
- 251000  
 NEUWEILER, G., F.P. MOHRES 1967  
 [DIE ROLLE DES ORTSGEDACHTNISSES BEI DER ORIENTIFRUNG DER  
 GROSSBLATT-FLEDERMAUSE, MEGADERMA LYRA]  
 Z. VERGL. PHYSIOL. 57, 147-171  
 GER ENG

- 240000  
ENG  
NEUWEILER, G., F.P. MOHRER 1967  
[ROLE OF SPACIAL MEMORY IN THE ORIENTATION]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL (ED.),  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 129-139  
200220  
ENG  
NEUWEILER, G., G. SCHULLER, H.U. SCHNITZLER 1971  
[ON-AND OFF-RESPONSES IN THE INFERIOR COLLICULUS OF THE GREATER  
HORSESHOE BAT TO PURE TONES]  
Z. VERGL. PHYSIOL. 74, 57-63  
221000  
ENG  
NORDMARK, J. 1960  
[PERCEPTION OF DISTANCE IN ANIMAL ECHOLOCATION]  
NATURE 188, 1009-1010  
263113  
ENG  
NOVICK, A. 1958  
[ORIENTATION OF PALEOTROPICAL BATS, I. MICROCHIROPTERA]  
J. EXP. ZOOL. 138, 81-154  
220103  
ENG  
NOVICK, A. 1962  
[ORIENTATION IN NEOTROPICAL BATS I. NATALIDAE AND EMBALLONURIDAE]  
J. MAMM. 43, 449-455  
220103  
ENG  
NOVICK, A. 1963  
[ORIENTATION IN NEOTROPICAL BATS II. PHYSILLOSTOMATIDAE AND  
DESMODONTIDAE]  
J. MAMM. 44, 44-56  
221000  
ENG  
NOVICK, A. 1963  
[PULSE DURATION IN THE ECHOLOCATION OF INSECTS BY THE BAT,  
PTERONOTUS]  
ERGER. DER BIOL. 26, 21-26  
221000  
ENG  
NOVICK, A. 1965  
[ECHOLOCATION OF FLYING INSECTS BY THE BAT, CHILONYCTERIS  
PSILOTTIS]  
BIOL. BULL. 128, 297-314  
221000  
ENG  
NOVICK, A. 1971  
[ECHOLOCATION IN BATS-SOME ASPECTS OF PULSE DESIGN]  
AMER. SCI. 59, 198-209

- 200110 ENG  
NOVICK, A., D.R. GRIFFIN 1961  
[LARYNGEAL MECHANISMS IN BATS FOR THE PRODUCTION OF ORIENTATION  
SOUNDS]  
J. EXP. ZOOL. 148, 125-145
- 221000 ENG  
NOVICK, A., J.R. VAISNYS 1964  
[ECHOLLOCATION OF FLYING INSECTS BY THE BAT, CHILONYCTERIS  
PARNELLII]  
BIOL. BULL. 127, 478-488
- 271313 ENG  
NOVICK, A., N. LEEN 1970  
[WORLD OF BATS]  
HOLT, REINHART, AND WINSTON, NEW YORK
- 240013 ENG  
NOVICK, A., R.A. DALE 1972  
[FORAGING BEHAVIOR IN FISHING BATS AND THEIR INSECTIVOROUS  
RELATIVES]  
J. MAMM. 52, 817-818
- 141 200310 ENG  
ORMEROD, F.C., J.D. PYE 1965  
[ECHOLLOCATION IN BATS]  
ACTA OTO-LARYNG. 53, 196-201 AD 611 302
- 212000 ENG  
PEFF, T.C., J.A. SIMMONS 1972  
[HORIZONTAL ANGLE RESOLUTION BY ECHOLLOCATING BATS]  
J. ACOUST. SOC. AM. 51, 2063-2065
- 200250 ENG  
POLLAK, G., O. W. HENSON, JR., A. NOVICK 1972  
[COCHLEAR MICROPHONIC AUDIOGRAMS IN THE [PURE TONE] BAT, CHILONYC-  
TERIS P. PARNELLII]  
SCIENCE 176, 66-68
- 200250 ENG  
POLLAK G. 1972  
[MIDDLE EAR MUSCLE ACTIVITY IN THE [PURE TONE] BAT, CHILONYC-  
TERIS P. PARNELLII]  
UNPUBLISHED MANUSCRIPT
- 220210 ENG  
PYE, A. 1966  
[STRUCTURE OF THE COCHLEA IN CHIROPTERA I. MICROCHIROPTERA-  
EMBALLONUROIDEA AND RHINOLOPHOIDEA]  
J. MORPH. 118, 495-510



220212 ENG  
 PYE, A. 1967  
 [STRUCTURE OF COCHLEA IN CHIROPTERA III. MICROCHIROPTERA-  
 PHYLLOSTOMATOIDEA]  
 J. MORPH. 121, 241-254

221010 ENG  
 PYE, J.D. 1960  
 [A THEORY OF ECHOLOCATION BY BATS]  
 J. LARNGYGOLOGY AND OTOTOLOGY 74, 718-729

221310 ENG  
 PYE, J.D. 1961  
 [ECHOLOCATION BY BATS]  
 ENDEAVOR 20, 101-111

221310 ENG  
 PYE, J.D. 1963  
 [ACTIVE ENERGY RADIATING SYSTEMS-THE BAT AND ULTRASONIC PRINCIPLES  
 I]  
 PROC. OF THE INTER. CONG. ON TECH. AND BLINDNESS, L.L. CLARK [ED.]  
 AMER. FOUND. FOR THE BLIND, NEW YORK, 35-49

203000 ENG  
 PYE, J.D. 1963  
 [MECHANISMS OF ECHOLOCATION]  
 ERGRN. DER BIOL. 26, 12-30

220110 ENG  
 PYE, J.D. 1967  
 [SYNTHESIZING THE WAVEFORMS OF BATS PULSES]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
 LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 43-67

220100 ENG  
 PYE, J.D. 1968  
 [ANIMAL SONAR IN AIR]  
 ULTRASONICS 6, 32-38

200210 ENG  
 PYE, J.D. 1968  
 [HEARING IN BATS]  
 SYMP. CIHA FOUND.-HEARING MECHANISMS IN VERTEBRATES, DE REUCK AND  
 KNIGHT [EDS.], 66-84

241201 ENG  
 PYE, J.D. 1971  
 [BATS AND FOG]  
 NATURE, 229, 57-574

- 220111  
 PYE, J. O. 1972  
 [BIMODAL DISTRIBUTION OF CONSTANT-FREQUENCIES IN SOME HIPPOSIDERID  
 BATS (MAMMALIA: HIPPOSIDERIDAE)]  
 J. ZOOL. LOND. 166, 323-335  
 ENG
- 240210  
 PYE, J.O., M. FLINN, A. PYE 1962  
 [CORRELATED ORIENTATION SOUNDS AND EAR MOVEMENTS OF HORSESHOE  
 BATS]  
 NATURE 196, 1185-1188  
 ENG
- 220210  
 PYE, J.O., L.H. ROBERTS 1970  
 [EAR MOVEMENTS IN A HIPPOSIDERID BAT]  
 NATURE 225, 285-286  
 ENG
- 220110  
 ROBERTS, L. H. 1972  
 [VARIABLE RESONANCE IN CONSTANT FREQUENCY BATS]  
 J. ZOOL. LOND. 166, 337-348  
 GER ENG
- 200002  
 ROER, H. 1969  
 [ZUR ERNAHRUNGSBIOLOGIE VON PLECOTUS AURITUS]  
 BONN. ZOOL. BEITR. 20, 378-383  
 ENG
- 200002  
 ROEDER, K.O. 1963  
 [ECHOES OF ULTRASONIC PULSES FROM FLYING MOTHS]  
 BIOL. BULL. 124, 200-209  
 ENG
- 231000  
 SCHNITZLER, H.U. 1967  
 [DISCRIMINATION OF THIN WIRES BY FLYING HORSESHOE BATS  
 (RHINOLOPHIDAE)]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
 LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 69-87  
 GER ENG
- 220100  
 SCHNITZLER, H.U. 1967  
 [KOMPENSATION VON DOPPLEREFFekten BEI HUFSEISEN-FLEDERMAUSEN. (COM-  
 PENSATION OF DOPPLER EFFECTS IN HORSESHOE BATS)]  
 DIE NATURWISS. 54, 523

231000 GER ENG  
 SCHNITZLER, H.U. 1968  
 [DIE ULTRASCHALL-ORTUNGSLAUTE DER HUFERSEN-FLEDERMAUSE  
 [CHIROPTERA-RHINOLOPHIDAE] IN VERSCHIEDENEN ORIENTIERUNGSSITUATIONEN]  
 Z. VERGL. PHYSIOL. 57, 376-408

220210 GER ENG  
 SCHNITZLER, H.U. 1970  
 [ECHOORTUNG REI DER FLEDERMAUS CHILONYCTERIS RUBIGINOSA]  
 Z. VERGL. PHYSIOL. 68, 25-38

220200 ENG  
 SCHNITZLER, H.U. 1970  
 [COMPARISON OF THE ECHOLOCATION BEHAVIOR IN RHINOLOPHUS FERRUM-EQUINUM AND CHILONYCTERIS RUBIGINOSA]  
 BIJDAGEN TOT DE DIERKUNDE. 40, 77-80

240000 GER ENG  
 SCHNITZLER, H.U. 1971  
 [FLEDERMAUSE IM WINDKANAL]  
 Z. VERGL. PHYSIOL. 73, 209-221

200220 GER ENG  
 SCHNITZLER, H.U., G. SCHULLER, G. NEUWEILER 1971  
 [ANTWORTEN DES COLLICULUS INFERIOR DER FLEDERMAUS RHINOLOPHUS EURYALE AUF TONALE REIZUNG]  
 DIE NATURWISS. 58, 627

200240 ENG  
 SCHULLER, G., G. NEUWEILER, H. U. SCHNITZLER 1971  
 [COLLICULAR RESPONSES TO THE FREQUENCY MODULATED FINAL PART OF ECHOLOCATION SOUNDS IN RHINOLOPHUS FERRUM EQUINUM]  
 Z. VERGL. PHYSIOL. 74, 153-155

220000 ENG  
 SIMKIN, G.N. 1969  
 [MOTION OF BATS TOWARD PLANE TARGETS AND THE NATURE OF CHANGES OF ECHOLOCATING SIGNALS IN THE PROCESS OF TARGET SELECTION AND RECOGNITION]  
 JPRS 48123 (28 MAY 1969)

262000 ENG  
 SIMKIN, G.N. 1970  
 [ECHOLOCATION PROCESS IN OXYGNATHOUS BATS IN FREE FLIGHT]  
 JPRS 50665 (4 JUNE 1970)

263000 RUS ENG  
SIMKIN, G.N. 1970  
[ECHOLLOCATION, MEANS OF RECOGNITION, AND DIRECTION FINDING IN  
SMOOTH-NOSED BATS]  
MOSCOW UNIV. VESTNIK, SERIYA 6, BIOLOGIYA, 25, 111-114  
260013 ENG  
SIMKIN, G.N., N.D. PATLYAKEVICH 1970  
[LOCATION SYSTEMS IN GENUS VESPERTILIO BATS AND FLITTERMICE]  
JPRS 51884 (1 DECEMBER 1970)  
273170 ENG  
SIMMONS, J.A. 1969  
[DEPTH PERCEPTION BY SONAR IN THE BAT]  
PRINCETON UNIV., DISSERTATION, UNIV. MICROFILMS, ANN ARBOR, MICH.  
220100 ENG  
SIMMONS, J.A. 1969  
[ACOUSTIC RADIATION PATTERNS FOR THE ECHOLLOCATING BATS CHILONYC-  
TERIS RUBIGINOSA AND EPTESICUS FUSCUS]  
J. ACOUS. SOC. AM. 46, 1054-1056  
233030 ENG  
SIMMONS, J.A. 1971  
[ECHOLLOCATION IN BATS-SIGNAL PROCESSING OF ECHOES FOR TARGET  
RANGE]  
SCIENCE 171, 925-928  
232040 ENG  
SIMMONS, J. A. 1971  
[THE SONAR RECEIVER OF THE BAT]  
ANN. N.Y. ACAD. SCI. 188, 161-174  
232040 ENG  
SIMMONS, J. A. 1972  
[RESPONSE OF THE DOPPLER SONAR SYSTEM IN THE BAT, RHINOLOPHUS  
FERRUM-EQUINUM]  
TO BE PUBLISHED IN J. ACOUST. SOC. AM.  
233030 ENG  
SIMMONS, J.A., J.A. VERNON 1971  
[ECHOLLOCATION-DISCRIMINATION OF TARGETS BY THE BAT, EPTESICUS  
FUSCUS]  
J. EXP. ZOOL. 176, 315-328

- 220110  
SOKOLOV, B.V., A.K. MAKAROV 1971  
[THE DIRECTION OF ULTRASONIC ORIENTATIONAL RADIATION OF THE LARGE  
RHINOLOPHIDAE AND THE ROLE OF NASAL OUTGROWTHS IN ITS FORMATION]  
SCIENTIFIC REPORTS OF THE HIGHER SCHOOL, BIOLOGICAL SCIENCES 7,  
37-44  
RUS ENG
- 223040  
STROTHER, G.K. 1961  
[NOTE ON THE POSSIBLE USE OF ULTRASONIC PULSE COMPRESSION BY BATS]  
J. ACOUST. SOC. AM. 33, 696-697  
ENG
- 200040  
STROTHER, G.K. 1967  
[COMMENTS ON AURAL PULSE COMPRESSION IN BATS AND HUMANS;  
J.J.G. MC CUE, J. ACOUST. SOC. AM. 40, 545-548, 1966]  
J. ACOUST. SOC. AM. 41, 529  
ENG
- 220110  
STROTHER, G.K., M. MOGUS 1970  
[ACOUSTICAL BEAM PATTERNS FOR BATS-SOME THEORETICAL  
CONSIDERATIONS]  
J. ACOUST. SOC. AM. 48, 1430-1432  
ENG
- 200260  
SUGA, N. 1964  
[RECOVERY CYCLES AND RESPONSES TO FREQUENCY MODULATED TONE  
PULSES IN AUDITORY NEURONS OF ECHO-LOCATING BATS]  
J. PHYSIOL. 175, 50-80  
ENG
- 200260  
SUGA, N. 1965  
[FUNCTIONAL PROPERTIES OF AUDITORY NEURONES IN THE CORTEX OF ECHO-  
LOCATING BATS]  
J. PHYSIOL. 181, 671-700  
ENG
- 200260  
SUGA, N. 1968  
[ANALYSIS OF FREQUENCY-MODULATED AND COMPLEX SOUNDS BY SINGLE  
AUDITORY NEURONES OF BATS]  
J. PHYSIOL. 198, 51-80  
ENG
- 231210  
SUGA, N. 1969  
[ECHO-LOCATION AND EVOKED POTENTIALS OF BATS AFTER ABLATION OF  
INFERIOR COLLICULUS]  
J. PHYSIOL. 203, 707-728  
ENG

231210 ENG  
 SUGA, N. 1969  
 [ECHO-LOCATION OF BATS AFTER ABLATION OF AUDITORY CORTEX]  
 J. PHYSIOL. 203, 729-739

200240 ENG  
 SUGA, N. 1970  
 [ECHO-RANGING NEURONS IN THE INFERIOR COLLICULUS OF BATS]  
 SCIENCE 170, 449-452

200240 ENG  
 SUGA, N. 1972  
 [NEURAL ATTENUATION OF RESPONSES TO EMITTED SOUNDS IN ECHOLocATING  
 BATS]  
 SCIENCE 177, 82-84

273000 ENG  
 SUTHERS, R.A. 1965  
 [ACOUSTIC ORIENTATION BY FISH CATCHING BATS]  
 J. EXPL. ZOOL. 158, 319-348

233000 ENG  
 SUTHERS, R.A. 1967  
 [COMPARATIVE ECHOLocATION BY FISHING BATS]  
 J. MAMM. 48, 79-87

240000 ENG  
 SUTHERS, R.A., J. CHASE 1969  
 [VISUAL FORM DISCRIMINATION BY ECHOLocATING BATS]  
 BIOL. BULL. 137, 535-546

200110 ENG  
 SUTHERS, R.A., S.P. THOMAS, B.J. SUTHERS 1972  
 [RESPIRATION, WING-BEAT AND ULTRASONIC PULSE EMISSION IN AN  
 ECHO-LOCATING BAT]  
 J. EXP. ZOOL. 56, 37-48

200240 ENG  
 VAN BERGEIJK, W.A. 1964  
 [SONIC PULSE-COMPRESSION IN BATS AND PEOPLE- A COMMENT]  
 J. ACOUST. SOC. AM. 36, 594-597

200250 RUS ENG  
 VASILYEV, A.G. 1967  
 [COMPARATIVE CHARACTERISTICS OF THE AUDITORY SYSTEM OF THE  
 VESPERTILIONID AND HORSESHOE BATS [ELECTROPHYSIOLOGICAL DATA]]  
 DAN SSSR 175, 1414-1417

- 221000  
 VERNON, J., E.A. PETERSON 1965  
 [ECHOLOCATION SIGNALS IN THE FREE-TAILED BAT, TADARIDA MEXICANA]  
 J. AUD. RES. 5, 317-330  
 ENG
- 200220  
 VERNON, J.A., J.I. DALLAND, E.G. WEVER 1966  
 [FURTHER STUDIES OF HEARING IN THE BAT, MYOTIS LUCIFUGUS, BY MEANS  
 OF COCHLEAR POTENTIALS]  
 J. AUD. RES. 6, 153-163  
 ENG
- 200230  
 VERNON, J., E. PETERSON 1966  
 [HEARING IN THE VAMPIRE BAT, DESMODUS ROTUNDUS MURINUS, AS SHOWN BY  
 COCHLEAR POTENTIALS]  
 J. AUD. RES. 6, 181-187  
 ENG
- 731330  
 VINCENT, F. 1964  
 [ACOUSTIC SIGNALS FOR AUTO-INFORMATION OR ECHOLOCACTION]  
 ACOUSTIC BEHAVIOR OF ANIMALS, R.G. BUSNEL [ED.],  
 ELSEVIER PUBL. CO., AMSTERDAM, 183-227  
 ENG
- 200013  
 WALKER, E.P. 1968  
 [MAMMALS OF THE WORLD]  
 THE JOHNS HOPKINS PRESS, BALTIMORE, VOL. 1, 182-392  
 ENG
- 263240  
 WEBSTER, F.A. 1963  
 [ACTIVE ENERGY RADIATING SYSTEMS-THE BAT AND ULTRASONIC  
 PRINCIPLES II, ACOUSTICAL CONTROL OF AIRBORNE INTERCEPTIONS BY  
 BATS]  
 PROC. OF THE INTER. CONG. ON TECH. AND BLINDNESS, L.L. CLARK [ED.]  
 AMER. FOUND. FOR THE BLIND, NEW YORK, 49-135  
 ENG
- 263000  
 WEBSTER, F.A. 1963  
 [BAT TYPE SIGNALS AND SOME IMPLICATIONS]  
 HUMAN FACTORS IN TECHNOLOGY, C. BENNETT [ED.], MC GRAW-HILL, NEW  
 YORK, 378-408  
 ENG
- 253000  
 WEBSTER, F.A. 1967  
 [INTERCEPTION PERFORMANCE OF ECHolocATING BATS IN THE PRESENCE  
 OF INTERFERENCE]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
 LAB. DE PHYSIOL. ACOUST. JOUY-EN-JOSAS, FRANCE, 673-713

240002

ENG

WEBSTER, F.A., D.R. GRIFFIN 1962  
[ROLE OF THE FLIGHT MEMBRANES IN INSECT CAPTURE BY BATS]  
ANIMAL BEHAVIOR 10, 332-340

273000

ENG

WEBSTER, F.A., O. G. BRAZIER 1965  
[EXPERIMENTAL STUDIES ON TARGET DETECTION, EVALUATION AND  
INTERCEPTION BY ECHOLOCATING BATS]  
AMRL TR 65 172, PP. 144 (NOVEMBER 1965) AD 628 055

273003

ENG

WEBSTER, F.A., O.G. BRAZIER 1968  
[EXPERIMENTAL STUDIES ON ECHOLOCATION MECHANISMS IN BATS]  
AMRL TR 67 192, PP. 165. (MAY 1968) AD 673 373

260003

ENG

WEBSTER, F.A., O.G. BRAZIER 1969  
[ECHOLOCATION INVESTIGATIONS ON BATS AND HUMANS-TARGET  
LOCALIZATION AND EVALUATION]  
AMRL TR 68 155, PP. 73, (SEPTEMBER 1969) AD 697 070

200210

RUS ENG

ZVORYKIN, V.P. 1959

[MORPHOLOGICAL BASIS OF LOCATIVE AND SUPERSONIC ABILITIES  
IN BAT]

ARKH. ANAT. GISTOL. I EMBRIOL. 36, 19-31



APPENDIX V

BIBLIOGRAPHY OF ACCUMULATED LITERATURE FOR  
WHALES, DOLPHINS, AND PORPOISES

**Preceding page blank**

LISTING FORMAT

		LANGUAGE	
6-DIGIT CODE		ORIGINAL	TRANSLATION
AUTHOR(S)	YEAR OF ISSUE		
TITLE OR DESCRIPTION			
SOURCE			

# KEY TO 6-DIGIT NUMBER CODE

## DIGITS

1	2	3	4	5	6
---	---	---	---	---	---

### DIGIT(S)

1

### FUNCTION(S)

#### ORDER

- 1 CETACEA
- 2 CHIROPTERA
- 4 OTHER

2-3

#### ECHOLOCATION

- 1 DETECTION
- 2 DISCRIMINATION
- 10 PERFORMANCE
- 20 SIGNALS ; FORMAT
- 40 BEHAVIOR

4-5

#### BIOLOGY

- 1 MORPHOLOGY ; ANATOMY
- 2 AUDIOMETRY
- 4 SIGNAL PROCESSING, CONDITIONING
- 10 SOUND GENERATION
- 20 SOUND RECEPTION

6

#### ECOLOGY

- 1 HABITAT
- 2 PREY

**Examples:**

- 231210 A report about bats (2). Includes measurements, description, or commentary about echolocation signals (e.g., characteristics, repetition rate) (20), and the bats' performance (10) on target or obstacle detection or avoidance (1) ( $20+10+1=31$ ), plus measurements, description or commentary on the bats' ability or capability to detect and/or localize sounds (echoes) (20), and the morphology and/or anatomy of the hearing organ (1) ( $20+1=21$ ). Does not include information about the bats' normal hunting habitat or food prey (0).
- 572101 A report about whales, dolphins, or porpoises (1), including measurements and/or commentary on other animal forms (4) ( $4+1=5$ ). Includes measurements, description, or commentary about the animals' echolocation signals (20), and their performance(s) (10) in discriminating between targets (2), plus commentary on the animals' behavior while echolocating (e.g., scanning motions, swimming speed) (40) ( $40+20+10+2=72$ ). Includes commentary on the mechanics of sound generation (e.g., beaming, energy requirements) (10), plus description of the animals' normal hunting habitat(s) (1).

132020

ENG

AIRAPETIANZ, E. SH., A. G. GOLUBKOV, I. V. YERSHOVA, A. R. ZHEZHERIN,  
V. I. KOROLEV 1969  
[ECHOLOCATION DIFFERENTIATION AND CHARACTERISTICS OF RADIATED  
PULSES IN DOLPHINS]  
DOKLADY AKADEMII NAUK SSSR 188, 1197-1199; JPRS 49479  
(19 DECEMBER 1969)

373313

RUS ENG

AYRAPET'YANTS, F. SH., A. I. KONSTANTINOV 1970  
[ECHOLOCATION IN NATURE]  
NAUKA PUBL. HOUSE, LENINGRAD. JPRS 51511  
(6 OCTOBER 1970) (CH. 17 AND CONCLUSION)

123040

ENG

ALTES, R. 1971  
[COMPUTER DERIVATION OF SOME DOLPHIN ECHOLOCATION SIGNALS]  
SCIENCE 173, 912-914

102200

ENG

ANDERSEN, S. 1970  
[DIRECTIONAL HEARING IN THE HARBOUR PORPOISE PHOCOENA PHOCOENA]  
INVESTIGATIONS ON CETACEA, G. PILLERI [ED.], VOL. 2,  
BENTELI AG, BERNE, 260-263

100220

ENG

ANDERSEN, S. 1970  
[AUDITORY SENSITIVITY OF THE HARBOUR PORPOISE PHOCOENA PHOCOENA]  
INVESTIGATIONS ON CETACEA, G. PILLERI [ED.], VOL. 2  
BENTELI AG, BERNE, 256-259

122040

ENG

APPLIED RESEARCH LABORATORIES,  
THE UNIVERSITY OF TEXAS AT AUSTIN, AND NAVAL UNDERSEA RESEARCH AND  
DEVELOPMENT CENTER, SAN DIEGO 1969  
[DELPHINID SONAR- PULSE WAVE AND SIMULATION STUDIES]  
NUC TP 175, PP. 83 (DECEMBER 1969)

120000

ENG

APPLIED RESEARCH LABORATORIES,  
THE UNIVERSITY OF TEXAS AT AUSTIN 1970  
[QUARTERLY STATUS REPORT NO. 3 UNDER CONTRACT N66001-70-C-0268  
FOR THE PERIOD 16 MARCH-15 JUNE 1970]  
ARL/UT, PP. 47 (3 AUGUST 1970)

Preceding page blank

- 120000  
APPLIED RESEARCH LABORATORIES,  
THE UNIVERSITY OF TEXAS AT AUSTIN 1971  
[QUARTERLY STATUS REPORT NO. 3 UNDER CONTRACT N66001-71-C-0369  
FOR THE PERIOD 25 MAY-24 AUGUST 1971]  
ARL/UT, PP. 18 (22 SEPTEMBER 1971)  
ENG
- 120000  
BACKUS, R.H., W.E. SCHEVILL 1966  
[P'YSETER CLICKS]  
WHALES, DOLPHINS, AND PORPOISES, K.S. NORRIS [ED.], UNIV. OF CALIF.  
PRESS, BERKELEY, 510-528  
ENG
- 112000  
BARTA, R.E. 1969  
[ACOUSTICAL PATTERN DISCRIMINATION BY AN ATLANTIC BOTTLENOSE  
DOLPHIN]  
NAVAL UNDERSEA CENTER, SAN DIEGO, CALIF., UNPUBLISHED REPORT, PP. 26  
RUS ENG
- 153000  
BAGDONAS, A., V.M. BEL#KOVICH, N.L. KRUSHINSKAYA 1970  
[INTERACTION OF ANALYZERS IN DOLPHINS DURING DISCRIMINATION OF  
GEOMETRICAL FIGURES UNDER WATER]  
J. HIGHER NEURAL ACT. 20, 1070-1075  
ENG
- 120040  
BATTEAU, D.W. 1967  
[DISCUSSION OF PAPER BY K.S. NORRIS]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 438-441  
ENG
- 100250  
BEL#KOVICH, V.M. 1970  
[ANATOMY AND FUNCTION OF THE EAR IN DOLPHINS]  
ZOOLOGICHESKII ZH. 2, 275-282; JPRS 50253 (7 APRIL 1970)  
ENG
- 140001  
BEL#KOVICH, V.M., N.L. KRUSHINSKAYA, V.S. GUREVICH 1969  
[THE BEHAVIOR OF DOLPHINS IN CAPTIVITY]  
PRIRODA, 18-28; JPRS 50701 (10 JUNE 1970)  
ENG
- 100010  
BLOOME, K.A. 1963  
[ON THE MICROANATOMY OF THE MELON AND SURROUNDING INTEGUMENT OF THE  
PORPOISE (TURSIOPS THUNCATUS)]  
LOCKHEED REPORT LR 17084, PP. 17 (21 OCTOBER 1963)

- 140310  
BOYNTON, K.L. 1966  
[THE REMARKABLE DOLPHIN]  
CANADIAN AUDUBON 28, 24-29
- 100270  
ENG  
BULLOCK, T.H., A.D. GRINNELL, E. IKEZONO, K. KAMEDA, Y. KATSUKI,  
M. NOMOTO, O. SATO, N. SUGA, K. YANAGISAWA 1968  
[ELECTROPHYSIOLOGICAL STUDIES OF CENTRAL AUDITORY MECHANISMS IN  
CETACEANS]  
Z. VERGL. PHYSIOL. 59, 117-156
- 100270  
ENG  
BULLOCK, T.H., S.H. RIDGWAY 1972  
[EVOKED POTENTIALS IN THE CENTRAL AUDITORY SYSTEM OF ALERT PORPOISES  
TO THEIR OWN AND ARTIFICIAL SOUNDS]  
J. NEUROBIOL. 3, 79-99
- 120000  
FRE  
BUSNEL, R.G., A. DZIEDZIC, S. ANDERSEN 1963  
[SUR CERTAINES CARACTERISTIQUES DES SIGNAUX ACOUSTIQUES DU  
MARSOULIN PHOCOENA (L.)]  
C.R. ACAD. SCI. PARIS 257, 2545-2548
- 113000  
FRE  
BUSNEL, R.G., A. DZIEDZIC, S. ANDERSEN 1965  
[SEUILS DE PERCEPTION DE SYSTEME SONAR DU MARSOULIN PHOCOENA  
PHOCOENA L., EN FONCTION DU DIAMETRE D'UN OBSTACLE FILLEFORME]  
C.R. ACAD. SCI. PARIS 260, 295-297
- 111000  
ENG  
BUSNEL, R.G., A. DZIEDZIC, S. ANDERSEN 1965  
[ROLE OF TARGET IMPEDANCE IN DETECTION THRESHOLD OF THE  
SONAR SYSTEM OF PHOCOENA PHOCOENA (PORPOISE)]  
PAPER PRESENTED AT SOCIETY OF BIOLOGY, PARIS 159, 69-74  
(13 JANUARY 1965)
- 120000  
ENG  
BUSNEL, R.G., A. DZIEDZIC 1966  
[ACOUSTIC SIGNALS OF THE PILOT WHALE GLOBICEPHALA MALAENA, AND OF  
THE PORPOISES DELPHINUS DELPHIS, AND PHOCOENA PHOCOENA]  
WHALES, DOLPHINS, AND PORPOISES, K.S. NORRIS (ED.),  
UNIV. OF CALIF. PRESS, BERKELEY, 607-646
- 120000  
FRE  
BUSNEL, R.G., A. DZIEDZIC, M.L. GALLIEN 1966  
[CARACTERISTIQUES PHYSIQUES DE CERTAINS SIGNAUX ACOUSTIQUES DE  
DELPHINIDE, STENO BREIDANENSIS L.]  
C.R. ACAD. SCI. PARIS 262, 143-146

131040

FRE

BUSNEL, R.G., A. DZIEDZIC 1967  
[RESULTATS METROLOGIQUES EXPERIMENTAUX DE L'ECHOLOCATION CHEZ LE  
PHOCAENA PHOCAENA, ET LEUR COMPARAISON AVEC CEUX DE CERTAINES  
CHAUVES-SOURIS]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 2, R.G. BUSNEL (ED.),  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 307-338

FRE

120000

BUSNEL, R.G., A. DZIEDZIC 1967  
[OBSERVATIONS SUR LE COMPORTEMENT ET LES EMISSIONS ACOUSTIQUES  
DU CACHALOT LORS DE LA CHASSE]  
BOCAGIANA 14, 1-15

FRE

120000

BUSNEL, R.G., A. DZIEDZIC 1968  
[CHARACTERISTIQUES PHYSIQUES DES SIGNAUX ACOUSTIQUES DE PSEUDORCA  
CRASSIDENS OWEN]  
EXTRAIT DE MAMMALIA 32, 1-5

FRE

120000

BUSNEL, R.G., A. DZIEDZIC 1968  
[ETUDE DES SIGNAUX ACOUSTIQUES ASSOCIES A DES SITUATIONS DE  
DETRESSE CHEZ CERTAINS CETACES ODONTOCETES]  
EXTRAIT DE ANNALES DE L'INSTITUT OCEANOGRAPHIQUE 46, 109-144

FRE

120010

BUSNEL, R.G., G. PILLERI, F.C. FRASER 1958  
[NOTES CONCERNANT LE DAUPHIN STENELLA STYX GRAY 1846]  
EXTRAIT DE MAMMALIA 32, 192-203

FRE

100210

BUSNEL, R.G., D. GIRAUD-SAUVEUR, M.C. LEGRAND, C. MARHIC,  
M. MILOCHE, M.-M. TRILLAT 1968  
[CARACTERES CRISTALLINS PARTICULIERS DES OSSELETS DE CETACES  
ODONTOCETES]

C.R. ACAD. SCI. PARIS 37, 865-868

FRE

120000

BUSNEL, R.G., A. DZIEDZIC, B. ESCUDIE 1969  
[AUTOCORRELATION ET ANALYSE SPECTRALE DES SIGNAUX [SONAR] DE DEUX  
ESPECES DE CETACES ODONTOCETES UTILISANT LES BASSES FREQUENCES]  
C.R. ACAD. SCI. PARIS 269, 365-367

FRE

120000

BUSNEL, R.G., B. ESCUDIE, A. DZIEDZIC, A. HELLION 1971  
[STRUCTURE DES CLICS DOUBLES D'ECHOLOCATION DU GLOBICEPHALE  
[CETACE ODONTOCETES]]  
C.R. ACAD. SCI. PARIS 272, 2459-2461



100210  
 CLARKE, R. 1948  
 [HEARING IN CETACEA]  
 NATURE 161, 979-980  
 ENG

120100  
 DIERCKS K.J., R.T. TROCHTA, C.F. GREENLAW, W.E. EVANS 1971  
 [RECORDING AND ANALYSIS OF DOLPHIN ECHOLOCATION SIGNALS]  
 J. ACOUST. SOC. AM. 49, 1729-1732  
 ENG

120040  
 DIERCKS, K.J., R.T. TROCHTA 1972  
 [ANIMAL SONAR: MEASUREMENTS AND MEANING]  
 J. ACOUST. SOC. AM. 51, 133(A)  
 ENG

102250  
 DREHER, J.J. 1967  
 [ACOUSTICAL HOLOGRAPHIC MODEL OF CETACEAN ECHOLOCATION]  
 ACOUSTICAL HOLOGRAPHY, VOL. 1, A.F. METHERELL, M.A. EL-SUM,  
 L. LARMORE [EDS.], PLENUM PRESS, NEW YORK, 127-137  
 ENG

120040  
 DREHER, J.J., W.W. SUTHERLAND, W.E. EVANS 1965  
 [INSTRUMENTATION TECHNIQUES FOR ANALYSIS OF CHARACTERISTICS OF  
 ANIMAL SONAR PULSES]  
 LOCKHEED-CALIF. CO. RPT. NO. 18525, PP. 43 (14 JANUARY 1965)  
 ENG

120000  
 DUBROVSKII, N.A., P.S. KRASNOV, A.A. TITOV 1971  
 [ON THE EMISSION OF ECHO-LOCATION SIGNALS BY THE AZOV SEA HARBOR  
 PORPOISE]  
 AKUST. ZH. 16, 521-525; SOVIET PHYSICS-ACOUSTICS 16, 444-447  
 RUS

132000  
 DUBROVSKY, N.A., A.A. TITOV, P.S. KRASNOV, V.P. BABKIN,  
 V.M. LEKOMTSEV, G.V. NIKOLENKO 1970  
 [INVESTIGATION OF THE PERMISSION (SIC) CAPACITY OF THE  
 BLACK SEA TURSIOPS TRUNCATUS ECHOLOCATION APPARATUS]  
 TRUDY AKUST. INST. 19, 163-181  
 ENG

102200  
 DUDOK VAN HEFL, W.H. 1959  
 [AUDIO-DIRECTION FINDING OF THE PORPOISE, PHOCAENA PHOCAENA]  
 NATURE 183, 1063  
 ENG

143211  
 DUDOK VAN HEEL, W.H. 1962  
 [SOUND AND CETACEA]  
 NETHERLANDS J. SEA RES. 1, 407-507

- 143001  
DUDOK VAN HEEL, W.H. 1966  
[NAVIGATION IN CETACEA]  
WHALES, DOLPHINS, AND POMPOISES, K.S. NORRIS [ED.],  
UNIV. OF CALIF. PRESS, HERKELEY, 597-606  
ENG
- 120001  
DUNN, J.L. 1969  
[SPERM WHALE ACOUSTIC CHARACTERISTIC MEASUREMENTS FROM ASWEPS  
AIRCRAFT]  
U.S. NAV. OCEANOGRAPHIC OFFICE WASHINGTON, D.C., INFORMAL RPT.  
1R69-14, PP. 12 (JANUARY 1969)  
ENG
- 121040  
DZIEDZIC, A. 1967  
[INTERPRETATION THEORIQUE DE CERTAINES DONNEES EXPERIMENTALES  
SUR L'ECHOLLOCATION]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, K.G. BUSNEL [ED.],  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 339-356  
FRE
- 331000  
DZIEDZIC, A. (DATE UNKNOWN)  
[QUELQUES PERFORMANCES DES SYSTEMES DE DETECTION PAR ECHOS DES  
CHAUVES-SOURIS ET DES DELPHINIDAE]  
REVUE D'ACOUSTIQUE 1, 23-28  
FRE
- 321350  
DZIEDZIC, A. 1971  
[BIOLOGICAL SONARS]  
THEORIE ET APPLICATIONS DE L'ACOUSTIQUE SOUS-MARINE, CH. 16, PP. 68  
(UNPUBLISHED)  
FRE ENG
- 120000  
EREPHARDT, R.F., W.E. EVANS 1962  
[SOUND ACTIVITY OF THE CALIFORNIA GRAY WHALE, ESCHRICHTIUS  
GLAUCUS]  
J. AUD. ENG. SOC. 10, 324-328  
ENG
- 173220  
EINHORN, R.N. 1967  
[DOLPHINS CHALLENGE THE DESIGNER]  
ELECTRONIC DESIGN 25, 49-64  
ENG
- 321040  
ESCUDIE, B., A. HELLION, A. DZIEDZIC 1971  
[QUELQUES RESULTATS DANS L'ETUDE DES SONARS BIOLOGIQUES AERIENS ET  
MARINS PAR TRAITEMENT DU SIGNAL ET ANALYSE SPECTRALE]  
PROC. TROISIEME COLLOQUE SUR LE TRAITEMENT DU SIGNAL ET SES  
APPLICATIONS, NICE, FRANCE, 1-5 JUNE 1971, 533-557  
FRE ENG

- 120000  
EVANS, W.E. 1967  
[VOCALIZATION AMONG MARINE MAMMALS]  
MARINE BIO-ACOUSTICS, VOL. 2, W.N. TAVOLGA [ED.], PERGAMON PRESS,  
NEW YORK, 159-186  
ENG
- 163000  
EVANS, W.E. 1967  
[DISCUSSION OF PAPER BY A. GRINNELL]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 495-503  
ENG
- 173111  
EVANS, W.E. 1972  
[A DISCUSSION OF ECHOLLOCATION BY CETACEANS BASED ON EXPERIMENTS WITH  
MARINE DELPHINIDS AND ONE SPECIES OF FRESH WATER DOLPHIN]  
(TO BE PUBLISHED IN J. ACOUST. SOC. AM.)  
ENG
- 100210  
EVANS, W.E., J.J. DREHER 1961  
[SOME CONSIDERATIONS OF HEARING IN CETACEA]  
LOCKHEED AIRCRAFT CORP., CALIF. DIV., ENG. RES. LAB.,  
LTM 50017, PP. 25 (20 FEBRUARY 1961)  
ENG
- 161000  
EVANS, W.E., J.J. DREHER 1962  
[OBSERVATIONS ON SCOUTING BEHAVIOR AND ASSOCIATED SOUND PRODUCTION  
BY THE PACIFIC BOTTLENOSE PORPOISE (TURSIOPS GILLI DALL)]  
BULL. SO. CALIF. ACAD. SCI. 61, 217-226  
ENG
- 120110  
EVANS, W.E., J.H. PRESCOTT, 1962  
[OBSERVATIONS OF THE SOUND PRODUCTION CAPABILITIES OF THE  
BOTTLENOSE PORPOISE-A STUDY OF WHISLES AND CLICKS]  
ZOOLOGICA 47, 121-128  
ENG
- 100110  
EVANS, W.E., W.W. SUTHERLAND, R.G. BEIL 1964  
[DIRECTIONAL CHARACTERISTICS OF DELPHINID SOUNDS]  
MARINE BIO-ACOUSTICS, W.N. TAVOLGA [ED.], PERGAMON PRESS,  
NEW YORK, 353-372  
ENG
- 100310  
EVANS, W.E., J.J. DREHER 1964  
[THE REMARKABLE MAN OF THE SEA]  
NAVAL RES. REVIEWS 17, 17-22

133020

ENG

EVANS, W.E., B.A. POWELL 1966  
[CURRENT CETACEAN RESEARCH-A REVIEW]  
AAIA SYMPOSIUM ON MODERN DEV. IN MARINE SCIENCES, WESTERN  
PERIODICALS CO., LOS ANGELES, CALIF., PP. 11 (21 APRIL 1966)

132000

ENG

EVANS, W.E., B.A. POWELL 1967  
[DISCRIMINATION OF DIFFERENT METALLIC PLATES BY AN ECHOLOCATING  
DELPHINID]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 363-383

173330

ENG

EVANS, W.E., E.C. EVANS III 1971  
[ECHOLOCATION OF AQUATIC MAMMALS BASED ON EXPERIMENTAL EVIDENCE FROM  
MARINE AND FRESH WATER CETACEANS]  
UNPUBLISHED MANUSCRIPT, NAVAL UNDERSEA RESEARCH AND DEVELOPMENT  
CENTER, PP. 64 (APRIL 1971)

120000

ENG

FISH, M.P., W. MOWBRAY 1962  
[PRODUCTION OF UNDERWATER SOUND BY THE WHITE WHALE OR BELUGA,  
DELPHINAPTERUS LEUCAS (PALLAS)]  
J. MAR. RES. 20, 149-162

100210

ENG

FRASER, F.C., P.E. PURVES 1959  
[HEARING IN WHALES]  
ENDEAVOR, 18, 94-98

100210

ENG

FRASER, F.C., P.E. PURVES 1960  
[ANATOMY AND FUNCTION OF THE CETACEAN EAR]  
PROC. ROYAL SOC. LOND. 152, 62-77

100210

FRE ENG

GIPAUD-SAUVEUR, D. 1969  
[BIOPHYSICAL RESEARCH ON THE OSSICLES OF CETACEANS]  
EXTRAIT DE MAMMALIA 33, 285-340

112000

RUS ENG

GUREVICH, V.S. 1969  
[ECHOLOCATION DISCRIMINATION OF GEOMETRIC FIGURES IN THE DOLPHIN  
DELPHINUS DELPHIS]  
MOSCOW, VESTNIK MOSKOVSKOGO UNIVERSITETA, BIOLOGIYA, POCHVOVEDENIYE  
3, 109-112; JPRS 49281 (21 NOVEMBER 1969)

- 100220  
ENG  
HALL, J.D., C.S. JOHNSON 1972  
[AUDITORY THRESHOLDS OF A KILLER WHALE ORCINUS ORCA LINNAEUS]  
J. ACOUST. SOC. AM. 51, 515-517
- 160011  
ENG  
HERALD, E.S., R.L. BROWNELL, JR., F.L. FRYE, E.J. MORRIS,  
W.E. EVANS, A.B. SCOTT, 1969  
[BLIND RIVER DOLPHIN-FIRST SIDE SWIMMING CETACEAN]  
SCIENCE 166, 1408-1410
- 100220  
ENG  
JACOBS, D.W., J.D. HALL 1972  
[AUDITORY THRESHOLDS OF A FRESHWATER DOLPHIN, INIA GEOFFRENSIS,  
BLAINVILLE]  
J. ACOUST. SOC. AM. 51, 530-533
- 140000  
ENG  
JOHANNESSEN, C.L., J.A. HARDER 1960  
[SUSTAINED SWIMMING SPEEDS OF DOLPHINS]  
SCIENCE 132, 1550-1551
- 100220  
ENG  
JOHNSON, C.S. 1967  
[SOUND DETECTION THRESHOLDS IN MARINE MAMMALS]  
MARINE BIO-ACOUSTICS, VOL. 2, W.N. TAVOLGA [ED.], PERGAMON PRESS,  
NEW YORK, 247-260
- 123000  
ENG  
JOHNSON, C.S. 1967  
[POSSIBLE USE OF PHASE INFORMATION IN TARGET DISCRIMINATION AND  
THE ROLE OF PULSE RATE IN PORPOISE ECHO RANGING]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND RIONICS, VOL. 1, R.G. BUSNEL [ED.],  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 384-398
- 100220  
ENG  
JOHNSON, C.S. 1968  
[RELATION BETWEEN ABSOLUTE THRESHOLD AND DURATION OF TONE PULSES  
IN THE BOTTLENOSED PORPOISE]  
J. ACOUST. SOC. AM. 43, 757-763
- 100260  
ENG  
JOHNSON, C.S. 1968  
[MASKED TONAL THRESHOLDS IN THE BOTTLENOSED PORPOISE]  
J. ACOUST. SOC. AM. 44, 965-967

100220                      ENG  
JOHNSON, C.S. 1970  
[AUDITORY MASKING OF ONE PURE TONE BY ANOTHER IN THE BOTTLENOSED  
PORPOISE]  
J. ACOUST. SOC. AM. 49, 1317-1318  
100220                      ENG  
KELLOGG, W.N. 1953  
[ULTRASONIC HEARING IN THE PORPOISE, TURSIOPS TRUNCATUS]  
J. COMP. PHYSIOL. PSYCHOL. 46, 446-450  
153000                      ENG  
KELLOGG, W.N. 1958  
[ECHO RANGING IN THE PORPOISE]  
SCIENCE 128, 981-988  
153000                      ENG  
KELLOGG, W.N. 1959  
[SIZE DISCRIMINATION BY REFLECTED SOUND IN A BOTTLENOSED  
PORPOISE]  
J. COMP. PHYSIOL. PSYCHOL. 52, 509-514  
153000                      ENG  
KELLOGG, W.N. 1959  
[AUDITORY PERCEPTION OF SUBMERGED OBJECTS BY PORPOISES]  
J. ACOUST. SOC. AM. 31, 1-6  
140000                      ENG  
KELLOGG, W.N. 1960  
[AUDITORY SCANNING IN THE DOLPHIN]  
PHYSIOL. RECORD 10, 25-27  
173330                      ENG  
KELLOGG, W.N. 1961  
[PORPOISES AND SONAR]  
UNIV. OF CHICAGO PRESS, CHICAGO  
100220                      ENG  
KELLOGG, W.N., R. KOHLER 1952  
[REACTIONS OF THE PORPOISE TO ULTRASONIC FREQUENCIES]  
SCIENCE 116, 25 -252  
120000                      ENG  
KELLOGG, W.N., R. KOHLER 1953  
[PORPOISE SOUNDS AS SONAR SIGNALS]  
SCIENCE 117, 233-243  
132002                      RUSS  
KONSTANTINOV, A.I., N.F. MELNIKOV, A.A. TITOV 1968  
[ON THE ABILITY OF DOLPHINS TO RECOGNIZE OBJECTS]  
TR. DOKL. II. RFSPUHL. KONF. PO BIONIKE, KIEV, 57-59

- 100010  
 LAWRENCE, B., W.E. SCHEVILL 1956  
 [THE FUNCTIONAL ANATOMY OF THE DELPHINID NOSE]  
 BULL. COMP. ZOOL. 114, 103-151  
 100013  
 LAYNE, J.N. 1958.  
 [OBSERVATIONS ON FRESHWATER DOLPHINS IN THE UPPER AMAZON]  
 J. MAMM. 39, 1-22  
 100011  
 LEATHERWOOD, S., W.E. EVANS, S.W. RICE 1972  
 [THE WHALES, DOLPHINS, AND PORPOISES OF THE EASTERN  
 NORTH PACIFIC: A GUIDE TO THEIR IDENTIFICATION IN THE  
 WATER]  
 NAVAL UNDERSEA CENTER TECH. PUBL. 282, PP. 184 (MARCH 1972)  
 120000  
 LEVENSON, C. 1970  
 [MEGAPTERA SOURCE LEVELS]  
 USN OCEANOGRAPHIC OFF., CORRESPONDENCE WITH W.E. EVANS,  
 NUC-SAN DIEGO, PP. 2  
 103040  
 LEVY, J.C. 1967  
 [TYPES AND PROTOCOLS OF EXPERIMENTS TO BE PERFORMED IN ORDER TO  
 OBTAIN COMPARATIVE RESULTS]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 2, R.G. BUSNEL (ED.),  
 LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 1139-1150  
 120000  
 LILLY, J.C., A.M. MILLER 1961  
 [SOUNDS EMITTED BY THE BOTTLENOSE DOLPHIN]  
 SCIENCE 133, 1689-1693  
 100230  
 MC CORMICK, J.G. 1968  
 [THEORY OF HEARING FOR DELPHINIDS]  
 DISSERTATION, PRINCETON UNIV., UNIV. MICROFILMS, ANN ARBOR, MICH.  
 100230  
 MC CORMICK, J.G., E.G. WEVER, J. PALIN, S.H. RIDGWAY 1970  
 [SOUND CONDUCTION IN THE DOLPHIN EAR]  
 J. ACOUST. SOC. AM. 48, 1418-1428  
 120000  
 MIZUE, K., A. TAKEMURA, K. NAKASAI 1968  
 [UNDERWATER SOUND OF THE CHINESE FINLESS PORPOISE CAUGHT IN THE  
 JAPANESE COASTAL SEA]  
 BULL. OF THE FAC. OF FISHERIES, NAGASAKI UNIV. 25, 25-32

120002

ENG

MIZUE, K., A. TAKEMURA, K. NAKASAI 1969  
[UNDERWATER SOUND OF THE FALSE KILLER WHALE]  
BULL. OF THE FAC. OF FISHERIES, NAGASAKI UNIV. 28, 19-29

120000

ENG

MOHL, B., S. ANDERSEN 1972  
[ECHOLOCATION: HIGH FREQUENCY COMPONENT IN THE CLICK OF THE  
HARBOR PORPOISE (PHOCOENA PH. L.)]  
ZOOLOGISK INSTITUT, ARHUS UNIV., DENMARK (UNPUBLISHED MANUSCRIPT)  
PP. 16

121000

ENG

MOROZOV, V.P., A.I. AKOPIAN, V.I. BURDIN, K.A. ZAYTSEVA,  
YU. A. SOKOVYKH 1972  
[REpetition RATE OF RANGING SIGNALS OF DOLPHINS AS A FUNCTION OF  
DISTANCE TO TARGET]  
BIOFIZIKA 17, 139-144; JPHS 55729 (17 APRIL 1972)

133310

ENG

NORRIS, K.S. 1964  
[SOME PROBLEMS OF ECHOLOCATION IN CETACEANS]  
MARINE BIO-AcouSTICS, W.N. TAVOLGA [ED.], PERGAMON PRESS,  
NEW YORK, 317-336

140310

ENG

NORRIS, K.S. 1968  
[Evolution OF Acoustic MECHANISMS IN ODONTOCETE CETACEANS]  
Evolution AND ENVIRONMENT, E.T. DRAKE, [ED.], YALE UNIV. PRESS  
NEW HAVEN, 297-324

173330

ENG

NORRIS, K.S. 1969  
[THE ECHOLOCATION OF MARINE MAMMALS]  
THE BIOLOGY OF MARINE MAMMALS, H.T. ANDERSEN [ED.], ACADEMIC PRESS,  
NEW YORK, 391-423

173000

ENG

NORRIS, K.S., J.H. PRESCOTT, P.V. ASA-DORIAN, P. PERKINS 1961  
[EXPERIMENTAL DEMONSTRATION OF ECHOLOCATION BEHAVIOR IN THE  
PORPOISE, TUHSIOPS THUNCATUS [MONTAGU]]  
BIOL. HULL. 20, 163-176

172040

ENG

NORRIS, K.S., W.E. EVANS, R.N. TURNER 1967  
[ECHOLOCATION IN AN ATLANTIC BOTTLENOSE PORPOISE DURING  
DISCRIMINATION]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 409-437



- 120140  
ENG  
NORRIS, K.S., W.E. EVANS 1967  
[DIRECTIONALITY OF ECHOLOCATION CLICKS IN THE ROUGH-TOOTH PORPOISE  
STENO BREDANENSIS (LESSON)]  
MARINE BIO-AcouSTICS, VOL. 2, W.N. TAVOLGA [ED.], PERGAMON PRESS,  
NEW YORK, 305-316
- 100110  
ENG  
NORRIS, K.S., K.J. DORMER, J. PEGG, G.J. LIESE 1971  
[MECHANISM OF SOUND PRODUCTION AND AIR RECYCLING IN PORPOISES-A  
PRELIMINARY REPORT]  
CONTRIBUTION NO. 90, THE OCEANIC INST., WAIMANALO, HA.,  
PP. 13 (UNPUBLISHED MANUSCRIPT)
- 100210  
ENG  
OGAWA, T., S. ARIFUKU 1948  
[ON THE ACOUSTIC SYSTEM IN THE CETACEAN BRAINS]  
SCI. RPTS. RES. INST., TOKYO, 2, PP. 20
- 171000  
ENG  
PENNER, R.H., A.E. MURCHISON 1970  
[EXPERIMENTALLY DEMONSTRATED ECHOLOCATION IN THE AMAZON RIVER  
PORPOISE INIA GEOFFRENSIS (BLAINVILLE)]  
NAVAL UNDERSEA RESEARCH AND DEVELOPMENT CENTER TECH. PUBL. 187,  
PP. 30 (JUNE 1970)
- 120000  
ENG  
PERKINS, P.J., M.P. FISH, W.M. MOWBRAY 1966  
[UNDERWATER COMMUNICATION SOUNDS OF THE SPERM WHALE, PHYSETER  
CATODON]  
NORSK HVALFANGST-TIDENDE 12, 225-228
- 140003  
ENG  
PILLERI, G. 1970  
[OBSERVATIONS ON THE BEHAVIOUR OF PLATANISTA GANGETICA  
IN THE INDUS AND BRAHMAPUTRA RIVERS]  
INVESTIGATIONS ON CETACEA, G. PILLERI [ED.], VOL. 2,  
BENTELI AG, BERNE, 27-60
- 120000  
ENG  
PILLERI, G., C. KRAUS, M. GIHR 1971  
[PHYSICAL ANALYSIS OF THE SOUNDS EMITTED BY PLATANISTA INDI]  
INVESTIGATIONS ON CETACEA, G. PILLERI [ED.] VOL. 3(1),  
BENTELI AG, HERNE, 22-30

- 160001  
 ENG  
 POULTER, T.C. 1968  
 [VOCALIZATION OF THE GRAY WHALES IN LAGUNA OJO DE LIEBRE  
 (SCAMMONS LAGOON) RAJA CALIFORNIA, MEXICO]  
 NORSK HVALFANGST- TIDENDE 3, 50-62  
 120110  
 ENG  
 PURVES, P.E. 1967  
 [ANATOMICAL AND EXPERIMENTAL OBSERVATIONS ON THE CETACEAN SONAR  
 SYSTEM]  
 ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1, R.G. BUSNEL [ED.],  
 LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS, FRANCE, 197-270  
 100210  
 ENG  
 REYSENHACH DE HAAN, F.W. 1957  
 [HEARING IN WHALES]  
 ACTA OTO-LARYNGOLOGICA, SUPPLEMENTUM 134, PP. 114  
 100210  
 ENG  
 REYSENHACH DE HAAN, F.W. 1966  
 [LISTENING UNDERWATER-THOUGHTS ON SOUND AND CETACEAN HEARING]  
 WHALES, DOLPHINS, AND PORPOISES, K.S. NORRIS [ED.], UNIV. OF CALIF.  
 PRESS, BERKELEY, 583-596  
 132040  
 RUS  
 REZNIK, A.M., V.M. SKURNYAKOV, A.G. CHUPAKOV 1970  
 [LOCATION ACTIVITY OF BLACK SEA TURSIOPS TRUNCATUS BEING  
 PRESENTED TARGETS]  
 TRUDY AKUST. INST. 12, 116-120  
 160003  
 ENG  
 RIDGWAY, S.M. 1966  
 [DALL PORPOISE, PHOCOENOIDES DALLI (TRUE)-OBSERVATIONS IN CAPTIVITY  
 AND AT SEA]  
 NORSK HVALFANGST-TIDENDE 5, 97-110  
 100220  
 ENG  
 RIDGWAY, S.M.  
 [STUDIES ON DIVING DEPTH AND DURATION IN TURSIOPS TRUNCATUS]  
 PROC. III ANN. CONF. BIOL. SONAR AND DIV. MAMM., 23-24 MAY 1966,  
 STAN. RES. INST., 151-157  
 133310  
 RUS ENG  
 ROMANENKO, E.V. 1964  
 [UNDERWATER ECHOLLOCATION (SONAR) CAPACITY OF DOLPHINS (REVIEW)]  
 AKUST. Zh. 10, 385-397; SOV. PHYS.-ACOUST. 10, 331-342

- 100110 RUS ENG  
ROMANENKO, E.V., A.G. TOMILIN, B.A. ARTEMENKO 1965  
[ON THE QUESTION OF SOUND FORMATION AND THE DIRECTING OF SOUNDS IN  
DOLPHINS]  
BIONIKA, NAUKA, MOSCOW, 269-273
- 120110 ENG  
SCHEVILL, W.E., 1964  
[UNDERWATER SOUNDS OF CETACEANS]  
MARINE BIO-AcouSTICS, W.N. TAVOLGA [ED.], PERGAMON PRESS,  
NEW YORK, 307-316
- 100200 ENG  
SCHEVILL, W.E., B. LAWRENCE 1953  
[HIGH FREQUENCY AUDITORY RESPONSE OF A BOTTLENOSED PORPOISE,  
TURSIOPS TRUNCATUS (MONTAGU)]  
J. ACOUST. SOC. AM. 25, 1016-1017
- 100200 ENG  
SCHEVILL, W.E., B. LAWRENCE 1953  
[AUDITORY RESPONSE OF A BOTTLENOSED PORPOISE TURSIOPS TRUNCATUS,  
TO FREQUENCIES ABOVE 100KC]  
J. EXP. ZOOL. 124, 147-165
- H-151000 ENG  
SCHEVILL, W.E., B. LAWRENCE 1956  
[FOOD FINDING BY A CAPTIVE PORPOISE (TURSIOPS TRUNCATUS)]  
BREVIOIRA (MUS. COMP. ZOOL., CAMBRIDGE, MASS.) 53, PP. 15
- 160000 ENG  
SCHEVILL, W.E., W.A. WATKINS, 1966  
[SOUND STRUCTURE AND DIRECTIONALITY IN ORCINUS (KILLER WHALE)]  
ZOOLOGICA 51, 71-76
- 120110 ENG  
SCHEVILL, W.E., W.A. WATKINS, C. RAY 1969  
[CLICK STRUCTURE IN THE PORPOISE, PHOCOENA PHOCOENA]  
J. MAMM. 50, 721-728
- 160000 ENG  
SCHEVILL, W.E., WATKINS, W.A. 1971  
[PULSED SOUNDS OF THE PORPOISE LAGENORHYNCHUS AUSTRALIS]  
BREVIOIRA (MUS. COMP. ZOOL., CAMBRIDGE, MASS.) 366, PP. 10
- 120040 ENG  
SINGLETON, R.C., I.C. POULTER 1967  
[SPECTRAL ANALYSIS OF THE CALL OF THE MALE KILLER WHALE]  
IEEE TRANS. ON AUDIO AND ELECTROACOUST. AU-15, 104-113

- 112011  
ENG  
SOKOLOV, V. 1971  
[CETACEAN RESEARCH IN THE USSR]  
INVESTIGATIONS ON CETACEA, G. PILLERI [ED.], VOL. 3(2),  
BENTELI AG, HERNE, 317-339
- 161210  
ENG  
SPONG, P., D. WHITE 1969  
[CETACEAN RESEARCH AT THE VANCOUVER PUBLIC AQUARIUM]  
TECH. RPT. NO. 2, UNIV. OF BRITISH COLUMBIA, VANCOUVER, B.C.  
DIV. OF NEUROLOGICAL SCIENCES, CETACEAN RES. LAB.,  
PP. 117 (SUMMER 1969)
- 100220  
RUS  
SUPIN, A. YA., M.N. SUKCHORUCHKO 1970  
[THE DETERMINATION OF AUDITORY THRESHOLDS IN PHOCOENA BY THE  
METHOD OF SKIN GALVANIC REACTION]  
TRUDY AKUST. INST. 12, 194-199
- 520310  
ENG  
TAVOLGA, W.N. 1965  
[REVIEW OF MARINE BIO-ACOUSTICS: STATE OF THE ART, 1964]  
TECH. RPT. DEPT. OF ANIMAL BEHAVIOR, AMER. MUS. NAT. HIST.,  
PP. 100 (FEBRUARY 1965)
- 171333  
ENG  
TOMILIN, A.G. 1966  
[ISTORIYA SLEPOGO KASHALOTA (HISTORY OF THE BLIND SPERM WHALE)]  
IZD. NAUKA, MOSCOW, 1965; ROUGH DRAFT TRANSLATION PREPARED BY  
TRANSLATION DIV., FOREIGN TECH. DIV., WP-AFB, OHIO, 1966-HF-66-434,  
PP. 208 (5 DECEMBER 1966)
- 153000  
ENG  
TURNER, R.N. 1964  
[METHODOLOGICAL PROBLEMS IN THE STUDY OF CETACEAN BEHAVIOR]  
MARINE BIO-ACOUSTICS, W.N. TAVOLGA [ED.], PERGAMON PRESS,  
NEW YORK, 337-352
- 172000  
ENG  
TURNER, R.N., K.S. NORRIS 1966  
[DISCRIMINATIVE ECHOCLOCATION IN A PORPOISE]  
J. EXPER. ANAL. BEHAV. 9, 535-544
- 100013  
ENG  
UNOKAWA, T., S. KITAMURA, Y. MOTOMURA, K. YAMAMOTO, T. KATAOKA  
1966  
[THE DOLPHINS OF MIE PREFECTURE]  
MIE SEIHUTSU (BIOLOGY OF MIE) 16, 49-52

120110

FRE

VINCENT, F. 1964

[ETUDES PRELIMINAIRES DE CERTAINES EMISSIONS ACOUSTIQUES DE DELPHINUS DELPHIS L. EN CAPTIVITE]

BULL. INST. OCEANOGR. MONACO, NO. 1172, PP. 23

731330

ENG

VINCENT, F. 1964

[ACOUSTIC SIGNALS FOR AUTO-INFORMATION OR ECHOLOCACTION]

ACOUSTIC BEHAVIOUR OF ANIMALS, R.G. BUSNEL [ED.],

ELSEVIER PUBL. CO., AMSTERDAM, 183-227

100013

ENG

WALKER, E.P. 1968

[MAMMALS OF THE WORLD]

THE JOHNS HOPKINS PRESS, BALTIMORE, VOL. 2, 1083-1145

100210

ENG

YAMADA, M. 1953

[CONTRIBUTION TO THE ANATOMY OF THE ORGAN OF HEARING OF WHALES]

SCI. REPORTS OF WHALES RES. INST., TOKYO 8, 1-79

100010

RUS ENG

ZVORYKIN, V.P. 1963

[MORPHOLOGICAL BASIS FOR THE ULTRASONIC AND SONIC DETECTION

CHARACTERISTICS OF THE DOLPHIN]

ARK. ANAT. GISTOL. I EMBRIOL. 7, 3-17; JPRS 21, 038

(11 SEPTEMBER 1963)

## **APPENDIX VI**

### **BIBLIOGRAPHY OF ACCUMULATED LITERATURE FOR OTHER ANIMALS**

**Preceding page blank**

## LISTING FORMAT

		LANGUAGE	
6-DIGIT CODE		ORIGINAL	TRANSLATION
AUTHOR(S)	YEAR OF ISSUE		
TITLE OR DESCRIPTION			
SOURCE			

# KEY TO 6-DIGIT NUMBER CODE

## DIGITS

1	2	3	4	5	6
---	---	---	---	---	---

### DIGIT(S)

1

### FUNCTION(S)

#### ORDER

- 1 CETACEA
- 2 CHIROPTERA
- 4 OTHER

2-3

#### ECHOLOCATION

- 1 DETECTION
- 2 DISCRIMINATION
- 10 PERFORMANCE
- 20 SIGNALS ; FORMAT
- 40 BEHAVIOR

4-5

#### BIOLOGY

- 1 MORPHOLOGY ; ANATOMY
- 2 AUDIOMETRY
- 4 SIGNAL PROCESSING, CONDITIONING
- 10 SOUND GENERATION
- 20 SOUND RECEPTION

6

#### ECOLOGY

- 1 HABITAT
- 2 PREY



Examples:

- 231210 A report about bats (2). Includes measurements, description, or commentary about echolocation signals (e.g., characteristics, repetition rate) (20), and the bats' performance (10) on target or obstacle detection or avoidance (1) ( $20+10+1=31$ ), plus measurements, description or commentary on the bats' ability or capability to detect and/or localize sounds (echoes) (20), and the morphology and/or anatomy of the hearing organ (1) ( $20+1=21$ ). Does not include information about the bats' normal hunting habitat or food prey (0).
- 572101 A report about whales, dolphins, or porpoises (1), including measurements and/or commentary on other animal forms (4) ( $4+1=5$ ). Includes measurements, description, or commentary about the animals' echolocation signals (20), and their performance(s) (10) in discriminating between targets (2), plus commentary on the animals' behavior while echolocating (e.g., scanning motions, swimming speed) (40) ( $40+20+10+2=72$ ). Includes commentary on the mechanics of sound generation (e.g., beaming, energy requirements) (10), plus description of the animals' normal hunting habitat(s) (1).

471000 EVANS, W.F., M.D. HAUGEN 1963  
[AN EXPERIMENTAL STUDY OF THE ECHOLOCATION ABILITY OF  
THE CALIFORNIA SEA LION ZALOPHUS CALIFORNIANUS (LESSON)]  
BULL. SO. CALIF. ACAD. SCI. 62, 165-175  
ENG

400200 GENTRY, K.L. 1967  
[UNDERWATER AUDITORY LOCALIZATION IN THE CALIFORNIA  
SEA LION (ZALOPHUS CALIFORNIANUS)]  
J. AUD. RES. 7, 187-193  
ENG

471000 GOULD, E., M.C. NEGUS, A. NOVICK 1964  
[EVIDENCE FOR ECHOLOCATION IN SHREWS]  
J. EXP. ZOOL. 156, 19-38  
ENG

431000 GRIFFIN, D.R., K.A. SUTHERS 1970  
[SENSITIVITY OF ECHOLOCATION IN CAVE SWIFTS]  
BIOLOGICAL BULL. 139, 495-501  
ENG

400230 MOHL, R. 1964  
[PRELIMINARY STUDIES ON HEARING IN SEALS]  
VIDENSK. MEDD. FRA DANSK NATURH. FOREN. 80, 127, 283-294  
ENG

400220 MOHL, R. 1964  
[FREQUENCY DISCRIMINATION IN THE COMMON SEAL AND A  
DISCUSSION OF THE CONCEPT OF UPPER HEARING LIMIT]  
UNDW. ACOUST. 2, 43-54  
ENG

400220 MOHL, R. 1964  
[AUDITORY SENSITIVITY OF THE COMMON SEAL IN AIR AND  
WATER]  
J. AUD. RES. 8, 27-38  
ENG

400250 NORRIS, A. 1964  
[PHYSICAL FACTORS IN DIRECTIONAL HEARING IN AEGOLIUS  
FUNKEUS (LINNE) (STIGIFORMES), WITH SPECIAL REFERENCE  
TO THE SIGNIFICANCE OF THE ASYMMETRY OF THE  
EXTERNAL EARS]  
ARKIV FOR ZOOLOGI 20, 181-204

460001	NOVICK, A. 1959 [ACOUSTIC ORIENTATION IN THE CAVE SWIFTLET] BIOL. HULL. 117, 497-503	ENG
453210	PAYNE, R.S. 1971 [ACOUSTIC LOCATION OF PREY BY BARN OWLS (TYTO ALBA)] J. EXP. BIOL. 54, 535-573; DIV. ENGIN. APPL. PHYS., HARVARD UNIV. TECH. RPT. NO. 1, PP. 105 (10 NOVEMBER 1961) AD 271, 031	ENG
431213	PAYNE, R.S., W.H. DHURY, JR. 1958 [MARKSMAN OF THE DARKNESS] NAT. HIST. 67, 316-323	ENG
420000	POULTER, T.C. 1963 [SONAR SIGNALS OF THE SEA LION] SCIENCE 139, 753-755	ENG
420000	POULTER, T.C. 1963 [THE SONAR OF THE SEA LION] IEEE TRANS. ULTRASONICS ENGIN. UE-10, 109-111	ENG
461000	POULTER, T.C. 1966 [THE USE OF ACTIVE SONAR BY THE CALIFORNIA SEA LION ZALOPHUS CALIFORNIANUS (LESSON)] J. AUD. HFS. 6, 165-173	ENG
450000	POULTER, T.C. 1968 [SEA LION ECHO RANGING] J. ACOUST. SOC. AM. 43, 1454(L)	ENG
471000	POULTER, T.C. 1969 [SONAR OF PENGUINS AND FUR SEALS] PHOC. CALIF. ACAD. SCI. 36, 363-380	ENG
433000	POULTER, T.C., JENNINGS, R.A. 1969 [SONAR DISCRIMINATION ABILITY OF THE CALIFORNIA SEA LION, ZALOPHUS CALIFORNIANUS] PHOC. CALIF. ACAD. SCI. 14, 381-389	ENG

450000  
SCHEVILL, W.F., 1968  
[SEA LION ECHO RANGING]  
J. ACOUST. SOC. AM. 43, 1458-1459(L)  
ENG

420000  
SCHEVILL, W.F., W.A. WATKINS, L. RAY 1963  
[UNDERWATER SOUNDS OF PINNIPEDS]  
SCIENCE 141, 50-53  
ENG

420000  
SCHEVILL, W.F., W.A. WATKINS 1965  
[UNDERWATER CALLS OF LEPTONYCHOTES (WEDDELL SEAL)]  
ZOOLOGICA 50, 45-46  
ENG

440000  
SCHUSTERMAN, R.J., 1965  
[VISIBILITY AND REINFORCEMENT AS VARIABLES  
INFLUENCING THE UNDERWATER CLICK VOCALIZATIONS  
OF A CALIFORNIA SEA LION]  
AMER. ZOOL. 5, 329  
ENG

473000  
SCHUSTERMAN, R.J., 1966  
[UNDERWATER CLICK VOCALIZATIONS BY A CALIFORNIA  
SEA LION : EFFECTS OF VISIBILITY]  
PSYCHOL. REC. 17, 129-136  
ENG

473000  
SCHUSTERMAN, R.J., 1967  
[PERCEPTION AND DETERMINANTS OF UNDERWATER VOCALIZATION  
IN THE CALIFORNIA SEA LION]  
ANIMAL SONAR SYSTEMS-BIOLOGY AND BIONICS, VOL. 1,  
R.G. BUSNELL (ED.), LAB. DE PHYSIOL. ACOUST., JOUY-EN-JOSAS,  
FRANCE, 535-617  
ENG

440000  
SCHUSTERMAN, R.J., R. GENTRY, J. SCHMOOK 1966  
[UNDERWATER VOCALIZATION BY SEA LIONS : SOCIAL  
AND MIRROR STIMULI]  
SCIENCE 154, 54-542  
ENG

460000  
SCHUSTERMAN, R.J., R. GENTRY, J. SCHMOOK 1967  
[UNDERWATER SOUND PRODUCTION BY CAPTIVE CALIFORNIA  
SEA LIONS, ZALOPHUS CALIFORNIANUS]  
ZOOLOGICA 52, 21-24  
ENG

420040

SHAYKH, H.N., I.C. PUULTER 1967  
[SEA LION ECHO HANGING]  
J. ACOUST. SOC. AM. 42, 428-427

ENG

# APPLIED RESEARCH LABORATORIES

